

# PHYS 1443 – Section 002

## Lecture #13

*Wednesday, Oct. 17, 2007*

*Dr. Jaehoon Yu*

- Potential Energy and the Conservative Force
  - Gravitational Potential Energy
  - Elastic Potential Energy
- Conservation of Energy
- Work Done by Non-Conservative Forces
- Energy Diagram and Equilibrium
- More General Gravitational Potential Energy



# Announcements

- 2<sup>nd</sup> term exam next Wednesday, Oct. 24
  - Covers from Ch. 4.1 to what we cover Monday, Oct. 22 (Ch 8 or 9)
  - Time: 1 – 2:20pm in class
  - Location: SH103
- Quiz results
  - Average: 3.9/8
    - Equivalent to 49/100
  - Top score: 8/8



**Physics Department**  
**The University of Texas at Arlington**  
**COLLOQUIUM**

---

**The Nuclear Equation of State: A Tool to  
Constrain In-Medium Hardronic  
Interactions and Gravitational Physics**

**Dr. Plamen Krastev**  
**Texas A&M University -Commerce**

**4:00 pm Wednesday, October 17, 2007**  
**Room 101 SH**


**Abstract**

One of the most important but also challenging problems in modern physics is to understand properties of matter under extreme conditions of density and pressure. These properties are reflected in what is known as the nuclear equation of state, which is the relationship between pressure and density (and some other state variables). The equation of state plays a key role in our understanding of numerous phenomena here on Earth, such as the dynamics of heavy-ion collisions, and also processes and objects with astrophysical significance, such as supernova explosions and neutron stars. In neutron star cores, it also provides a direct link between the strong interaction physics and general relativity within which models of neutron stars are constructed and their properties studied. After briefly reviewing our previous work on the equation of state of dense matter, I will focus on our most recent results on its applications to properties and structure of both static and (rapidly) rotating neutron stars, and constraining possible time variations of the gravitational constant  $G$ .

**Refreshments will be served in the Physics Library at 3:30 pm**

# Gravitational Potential Energy

*The potential energy given to an object by the gravitational field in the system of Earth due to the object's height from the surface*



*When an object is falling, the gravitational force,  $Mg$ , performs the work on the object, increasing the object's kinetic energy. So the potential energy of an object at a height  $y$ , which is the potential to do work is expressed as*

$$U_g = \vec{F}_g \cdot \vec{y} = |\vec{F}_g| |\vec{y}| \cos \theta = |\vec{F}_g| |\vec{y}| = mgy \quad U_g \equiv mgy$$

*The work done on the object by the gravitational force as the brick drops from  $y_i$  to  $y_f$  is:*

$$\begin{aligned} W_g &= U_i - U_f \\ &= mgy_i - mgy_f = -\Delta U_g \end{aligned}$$


*What does this mean?*

*Work by the gravitational force as the brick drops from  $y_i$  to  $y_f$  is the negative change of the system's potential energy*

**→ Potential energy was lost in order for the gravitational force to increase the brick's kinetic energy.**

# Conservative and Non-conservative Forces

*The work done on an object by the gravitational force does not depend on the object's path in the absence of a retardation force.*



When directly falls, the work done on the object by the gravitation force is  $W_g = mgh$

When sliding down the hill of length  $l$ , the work is

$$W_g = F_{g-\text{incline}} \times l = mg \sin \theta \times l \\ = mg(l \sin \theta) = mgh$$

How about if we lengthen the incline by a factor of 2, keeping the height the same??

Still the same amount of work 😊

$$W_g = mgh$$

So the work done by the gravitational force on an object is independent of the path of the object's movements. It only depends on the difference of the object's initial and final position in the direction of the force.

Forces like gravitational and elastic forces are called the conservative force

1. If the work performed by the force does not depend on the path.
2. If the work performed on a closed path is 0.

*Total mechanical energy is conserved!!*

$$E_M \equiv KE_i + PE_i = KE_f + PE_f$$

Wednesday, Oct. 17, 2007



PHYS 1443-002, Fall 2007

Dr. Jaehoon Yu

3

# Conservative Forces and Potential Energy

The work done on an object by a conservative force is equal to the decrease in the potential energy of the system

$$W_c = \int_{x_i}^{x_f} F_x dx = -\Delta U$$

*What does this statement tell you?*

The work done by a conservative force is equal to the negative change of the potential energy associated with that force.

*Only the changes in potential energy of a system is physically meaningful!!*

*We can rewrite the above equation in terms of the potential energy  $\mathcal{U}$*

$$\Delta U = U_f - U_i = -\int_{x_i}^{x_f} F_x dx$$

So the potential energy associated with a conservative force at any given position becomes

$$U_f(x) = -\int_{x_i}^{x_f} F_x dx + U_i$$

Potential energy function


*What can you tell from the potential energy function above?*

*Since  $\mathcal{U}_i$  is a constant, it only shifts the resulting  $\mathcal{U}_f(x)$  by a constant amount. One can always change the initial potential so that  $\mathcal{U}_i$  can be 0.*



# Example for Potential Energy

A bowler drops bowling ball of mass 7kg on his toe. Choosing the floor level as  $y=0$ , estimate the total work done on the ball by the gravitational force as the ball falls on the toe.



Let's assume the top of the toe is 0.03m from the floor and the hand was 0.5m above the floor.

$$U_i = mgy_i = 7 \times 9.8 \times 0.5 = 34.3J \quad U_f = mgy_f = 7 \times 9.8 \times 0.03 = 2.06J$$

$$W_g = -\Delta U = -(U_f - U_i) = 32.24J \cong 30J$$

b) Perform the same calculation using the top of the bowler's head as the origin.

What has to change?

First we must re-compute the positions of the ball in his hand and on his toe.

Assuming the bowler's height is 1.8m, the ball's original position is -1.3m, and the toe is at -1.77m.

$$U_i = mgy_i = 7 \times 9.8 \times (-1.3) = -89.2J \quad U_f = mgy_f = 7 \times 9.8 \times (-1.77) = -121.4J$$

$$W_g = -\Delta U = -(U_f - U_i) = 32.2J \cong 30J$$

# Elastic Potential Energy

*Potential energy given to an object by a spring or an object with elasticity in the system that consists of an object and the spring.*

*The force spring exerts on an object when it is distorted from its equilibrium by a distance  $x$  is*

$$F_s = -kx \quad \text{Hooke's Law}$$

*The work performed on the object by the spring is*

$$W_s = \int_{x_i}^{x_f} (-kx) dx = \left[ -\frac{1}{2}kx^2 \right]_{x_i}^{x_f} = -\frac{1}{2}kx_f^2 + \frac{1}{2}kx_i^2 = \frac{1}{2}kx_i^2 - \frac{1}{2}kx_f^2$$

*The potential energy of this system is*

$$U_s \equiv \frac{1}{2}kx^2$$

*What do you see from the above equations?*




*The work done on the object by the spring depends only on the initial and final position of the distorted spring.*

*Where else did you see this trend?*

*The gravitational potential energy,  $U_g$*

*So what does this tell you about the elastic force?*

*A conservative force!!!*






# Conservation of Mechanical Energy

*Total mechanical energy is the sum of kinetic and potential energies*

$$E \equiv K + U$$



Let's consider a brick of mass  $m$  at the height  $h$  from the ground

*What is the brick's potential energy?*

$$U_g = mgh$$

*What happens to the energy as the brick falls to the ground?*

$$\Delta U = U_f - U_i = -\int_{x_i}^{x_f} F_x dx$$

*The brick gains speed*

*By how much?*

$$v = gt$$

*So what?*

*The brick's kinetic energy increased*

$$K = \frac{1}{2}mv^2 = \frac{1}{2}mg^2t^2$$

*And?*

*The lost potential energy is converted to kinetic energy!!*

What does this mean?

*The total mechanical energy of a system remains constant in any isolated system of objects that interacts only through conservative forces:*


*Principle of mechanical energy conservation*

$$E_i = E_f$$

$$K_i + \sum U_i = K_f + \sum U_f$$

# Example

A ball of mass  $m$  at rest is dropped from the height  $h$  above the ground. a) Neglecting air resistance determine the speed of the ball when it is at the height  $y$  above the ground.



Using the principle of mechanical energy conservation

$$K_i + U_i = K_f + U_f \quad 0 + mgh = \frac{1}{2}mv^2 + mgy$$

$$\frac{1}{2}mv^2 = mg(h - y)$$

$$\therefore v = \sqrt{2g(h - y)}$$

b) Determine the speed of the ball at  $y$  if it had initial speed  $v_i$  at the time of the release at the original height  $h$ .

Again using the principle of mechanical energy conservation but with non-zero initial kinetic energy!!!

This result look very similar to a kinematic expression, doesn't it? Which one is it?

$$K_i + U_i = K_f + U_f$$


$$\frac{1}{2}mv_i^2 + mgh = \frac{1}{2}mv_f^2 + mgy$$

$$\frac{1}{2}m(v_f^2 - v_i^2) = mg(h - y)$$

$$\therefore v_f = \sqrt{v_i^2 + 2g(h - y)}$$

# Example

A ball of mass  $m$  is attached to a light cord of length  $L$ , making up a pendulum. The ball is released from rest when the cord makes an angle  $\theta_A$  with the vertical, and the pivoting point P is frictionless. Find the speed of the ball when it is at the lowest point, B.



Compute the potential energy at the maximum height,  $h$ . Remember where the 0 is.

$$h = L - L \cos \theta_A = L(1 - \cos \theta_A)$$

$$U_i = mgh = mgL(1 - \cos \theta_A)$$

Using the principle of mechanical energy conservation

$$K_i + U_i = K_f + U_f$$

$$0 + mgh = mgL(1 - \cos \theta_A) = \frac{1}{2}mv^2$$

$$v^2 = 2gL(1 - \cos \theta_A) \quad \therefore v = \sqrt{2gL(1 - \cos \theta_A)}$$

b) Determine tension  $T$  at the point B.

Using Newton's 2<sup>nd</sup> law of motion and recalling the centripetal acceleration of a circular motion

$$\begin{aligned} \sum F_r &= T - mg = ma_r = m \frac{v^2}{r} = m \frac{v^2}{L} \\ T &= mg + m \frac{v^2}{L} = m \left( g + \frac{v^2}{L} \right) = m \left( g + \frac{2gL(1 - \cos \theta_A)}{L} \right) \\ &= m \frac{gL + 2gL(1 - \cos \theta_A)}{L} \end{aligned}$$

$$\therefore T = mg(3 - 2\cos \theta_A)$$

Cross check the result in a simple situation. What happens when the initial angle  $\theta_A$  is 0?  $T = mg$

# Work Done by Non-conservative Forces

Mechanical energy of a system is not conserved when any one of the forces in the system is a non-conservative (dissipative) force.

Two kinds of non-conservative forces:

*Applied forces: Forces that are external to the system. These forces can take away or add energy to the system. So the mechanical energy of the system is no longer conserved.*

*If you were to hit a free falling ball, the force you apply to the ball is external to the system of ball and the Earth.*

$$W_{you} + W_g = \Delta K; \quad W_g = -\Delta U$$

*Therefore, you add kinetic energy to the ball-Earth system.*

$$W_{you} = W_{applied} = \Delta K + \Delta U$$

*Kinetic Friction: Internal non-conservative force that causes irreversible transformation of energy. The friction force causes the kinetic and potential energy to transfer to internal energy*

$$W_{friction} = \Delta K_{friction} = -f_k d$$

$$\Delta E = E_f - E_i = \Delta K + \Delta U = -f_k d$$



# Example of Non-Conservative Force

A skier starts from rest at the top of frictionless hill whose vertical height is  $20.0\text{m}$  and the inclination angle is  $20^\circ$ . Determine how far the skier can get on the snow at the bottom of the hill with a coefficient of kinetic friction between the ski and the snow is  $0.210$ .



Don't we need to know the mass?

Compute the speed at the bottom of the hill, using the mechanical energy conservation on the hill before friction starts working at the bottom

$$ME = mgh = \frac{1}{2}mv^2$$

$$v = \sqrt{2gh}$$

$$v = \sqrt{2 \times 9.8 \times 20.0} = 19.8\text{m/s}$$

The change of kinetic energy is the same as the work done by the kinetic friction.

What does this mean in this problem?

$$\Delta K = K_f - K_i = -f_k d$$

Since  $K_f = 0$   $-K_i = -f_k d; f_k d = K_i$

$$f_k = \mu_k n = \mu_k mg$$

$$d = \frac{K_i}{\mu_k mg} = \frac{\frac{1}{2}mv^2}{\mu_k mg} = \frac{v^2}{2\mu_k g} = \frac{(19.8)^2}{2 \times 0.210 \times 9.80} = 95.2\text{m}$$

Since we are interested in the distance the skier can get to before stopping, the friction must do as much work as the available kinetic energy to take it all away.

Well, it turns out we don't need to know the mass.

What does this mean?

No matter how heavy the skier is he will get as far as anyone else has gotten starting from the same height.