

PHYS 1443 – Section 002

Lecture #16

Wednesday, Oct. 31, 2007

Dr. Jae Yu

- Two Dimensional Collisions
- Center of Mass
- Fundamentals of Rotational Motions
- Rotational Kinematics
- Relationship between angular and linear quantities



Announcements

- Make sure you come to the colloquium today
 - 4pm in SH101
 - Refreshment at 3:30pm in SH108
- A midterm grade discussion next Monday, Nov. 5
 - Be sure not to miss this



**Physics Department
The University of Texas at Arlington
COLLOQUIUM**

**Exploring the Universe with
Balloon-borne Experiments**

Dr. Eun-Suk Seo
Institute for Physical Science & Technology
University of Maryland

4:00 pm Wednesday, October 31, 2007
Room 101 SH


Abstract

Cosmic rays, energetic particles coming from outer space, bring us information about the physical processes that accelerate particles to relativistic energies, about the effects of those particles in driving dynamical processes in our Galaxy, and about the distribution of matter and fields in interstellar space. The energy of these cosmic beams far exceeds energies produced by particle accelerators on Earth. Balloon-borne experiments are currently being used for understanding cosmic ray origin, acceleration and propagation, exploring the supernova acceleration limit, and searching for exotic sources such as dark matter and antimatter. Our on-going effort at Maryland with Balloon-borne Experiments will be presented and challenges of extending precision measurements to highest energy practical will be discussed.

Refreshments will be served in the Physics Library at 3:30 pm

Two dimensional Collisions

In two dimension, one needs to use components of momentum and apply momentum conservation to solve physical problems.



$$m_1 \vec{v}_{1i} + m_2 \vec{v}_{2i} = m_1 \vec{v}_{1f} + m_2 \vec{v}_{2f}$$

x-comp. $m_1 v_{1ix} + m_2 v_{2ix} = m_1 v_{1fx} + m_2 v_{2fx}$

y-comp. $m_1 v_{1iy} + m_2 v_{2iy} = m_1 v_{1fy} + m_2 v_{2fy}$

Consider a system of two particle collisions and scatters in two dimension as shown in the picture. (This is the case at fixed target accelerator experiments.) The momentum conservation tells us:

$$m_1 \vec{v}_{1i} + m_2 \vec{v}_{2i} = m_1 \vec{v}_{1i}$$

$$m_1 v_{1ix} = m_1 v_{1fx} + m_2 v_{2fx} = m_1 v_{1f} \cos \theta + m_2 v_{2f} \cos \phi$$

$$m_1 v_{1iy} = 0 = m_1 v_{1fy} + m_2 v_{2fy} = m_1 v_{1f} \sin \theta - m_2 v_{2f} \sin \phi$$

And for the elastic collisions, the kinetic energy is conserved:

Wednesday, Oct. 31, 2007




PHYS 1443-002, Fall 2007
Dr. Jaehoon Yu

What do you think we can learn from these relationships?

Example for Two Dimensional Collisions

Proton #1 with a speed 3.50×10^5 m/s collides elastically with proton #2 initially at rest. After the collision, proton #1 moves at an angle of 37° to the horizontal axis and proton #2 deflects at an angle ϕ to the same axis. Find the final speeds of the two protons and the scattering angle of proton #2, ϕ .



Since both the particles are protons $m_1 = m_2 = m_p$.

Using momentum conservation, one obtains

x-comp. $m_p v_{1i} = m_p v_{1f} \cos \theta + m_p v_{2f} \cos \phi$

y-comp. $m_p v_{1f} \sin \theta - m_p v_{2f} \sin \phi = 0$

Canceling m_p and putting in all known quantities, one obtains

$$v_{1f} \cos 37^\circ + v_{2f} \cos \phi = 3.50 \times 10^5 \quad (1)$$

$$v_{1f} \sin 37^\circ = v_{2f} \sin \phi \quad (2)$$

From kinetic energy conservation:

$$(3.50 \times 10^5)^2 = v_{1f}^2 + v_{2f}^2 \quad (3)$$

Solving Eqs. 1-3 equations, one gets

$$v_{1f} = 2.80 \times 10^5 \text{ m/s}$$

$$v_{2f} = 2.11 \times 10^5 \text{ m/s}$$

Do this at home 😊


Center of Mass

We've been solving physical problems treating objects as sizeless points with masses, but in realistic situations objects have shapes with masses distributed throughout the body.

Center of mass of a system is the average position of the system's mass and represents the motion of the system as if all the mass is on the point.

What does above statement tell you concerning the forces being exerted on the system?

The total external force exerted on the system of total mass M causes the center of mass to move at an acceleration given by $\vec{a} = \sum \vec{F} / M$ as if all the mass of the system is concentrated on the center of mass.




Consider a massless rod with two balls attached at either end.

The position of the center of mass of this system is the mass averaged position of the system

$$x_{CM} \equiv \frac{m_1 x_1 + m_2 x_2}{m_1 + m_2}$$

CM is closer to the heavier object

Motion of a Diver and the Center of Mass



(a)

Diver performs a simple dive.
The motion of the center of mass follows a parabola since it is a projectile motion.




(b)

Diver performs a complicated dive.
The motion of the center of mass still follows the same parabola since it still is a projectile motion.

The motion of the center of mass of the diver is always the same.

Example 9-12

Three people of roughly equivalent mass M on a lightweight (air-filled) banana boat sit along the x axis at positions $x_1=1.0\text{m}$, $x_2=5.0\text{m}$, and $x_3=6.0\text{m}$. Find the position of CM.




Using the formula
for CM

$$x_{CM} = \frac{\sum_i m_i x_i}{\sum_i m_i}$$

$$= \frac{M \cdot 1.0 + M \cdot 5.0 + M \cdot 6.0}{M + M + M} = \frac{12.0M}{3M} = 4.0(m)$$

Example for Center of Mass in 2-D

A system consists of three particles as shown in the figure. Find the position of the center of mass of this system.



Using the formula for CM for each position vector component

$$x_{CM} = \frac{\sum_i m_i x_i}{\sum_i m_i} \quad y_{CM} = \frac{\sum_i m_i y_i}{\sum_i m_i}$$

One obtains $\vec{r}_{CM} = x_{CM} \vec{i} + y_{CM} \vec{j} = \frac{(m_2 + 2m_3) \vec{i} + 2m_1 \vec{j}}{m_1 + m_2 + m_3}$

$$x_{CM} = \frac{\sum_i m_i x_i}{\sum_i m_i} = \frac{m_1 x_1 + m_2 x_2 + m_3 x_3}{m_1 + m_2 + m_3} = \frac{m_2 + 2m_3}{m_1 + m_2 + m_3}$$

If $m_1 = 2\text{kg}; m_2 = m_3 = 1\text{kg}$

$$y_{CM} = \frac{\sum_i m_i y_i}{\sum_i m_i} = \frac{m_1 y_1 + m_2 y_2 + m_3 y_3}{m_1 + m_2 + m_3} = \frac{2m_1}{m_1 + m_2 + m_3}$$

$$\vec{r}_{CM} = \frac{3\vec{i} + 4\vec{j}}{4} = 0.75\vec{i} + \vec{j}$$

Wednesday, Oct. 31, 2007



PHYS 1443-002, Fall 2007
Dr. Jaehoon Yu

Center of Mass of a Rigid Object

The formula for CM can be extended to a system of many particles or a Rigid Object

$$x_{CM} = \frac{m_1 x_1 + m_2 x_2 + \cdots + m_n x_n}{m_1 + m_2 + \cdots + m_n} = \frac{\sum_i m_i x_i}{\sum_i m_i}$$

$$y_{CM} = \frac{\sum_i m_i y_i}{\sum_i m_i}$$

$$z_{CM} = \frac{\sum_i m_i z_i}{\sum_i m_i}$$

The position vector of the center of mass of a many particle system is


$$\vec{r}_{CM} = x_{CM} \vec{i} + y_{CM} \vec{j} + z_{CM} \vec{k} = \frac{\sum_i m_i x_i \vec{i} + \sum_i m_i y_i \vec{j} + \sum_i m_i z_i \vec{k}}{\sum_i m_i}$$

$$\vec{r}_{CM} = \frac{\sum_i m_i \vec{r}_i}{M}$$

$$x_{CM} \approx \frac{\sum_i \Delta m_i x_i}{M}$$

$$x_{CM} = \lim_{\Delta m_i \rightarrow 0} \frac{\sum_i \Delta m_i x_i}{M} = \frac{1}{M} \int x dm$$


$$\vec{r}_{CM} = \frac{1}{M} \int \vec{r} dm$$



A rigid body – an object with shape and size with mass spread throughout the body, ordinary objects – can be considered as a group of particles with mass m_i densely spread throughout the given shape of the object

Example of Center of Mass; Rigid Body

Show that the center of mass of a rod of mass $\mathcal{M}$ and length $\mathcal{L}$ lies in midway between its ends, assuming the rod has a uniform mass per unit length.



The formula for CM of a continuous object is

$$x_{CM} = \frac{1}{M} \int_{x=0}^{x=L} x dm$$

Since the density of the rod (λ) is constant; $\lambda = M / L$

The mass of a small segment $dm = \lambda dx$

Therefore

$$x_{CM} = \frac{1}{M} \int_{x=0}^{x=L} \lambda x dx = \frac{1}{M} \left[\frac{1}{2} \lambda x^2 \right]_{x=0}^{x=L} = \frac{1}{M} \left(\frac{1}{2} \lambda L^2 \right) = \frac{1}{M} \left(\frac{1}{2} ML \right) = \frac{L}{2}$$

Find the CM when the density of the rod non-uniform but varies linearly as a function of x , $\lambda = \alpha x$

$$\begin{aligned} M &= \int_{x=0}^{x=L} \lambda dx = \int_{x=0}^{x=L} \alpha x dx \\ &= \left[\frac{1}{2} \alpha x^2 \right]_{x=0}^{x=L} = \frac{1}{2} \alpha L^2 \end{aligned}$$

$$x_{CM} = \frac{1}{M} \int_{x=0}^{x=L} \lambda x dx = \frac{1}{M} \int_{x=0}^{x=L} \alpha x^2 dx = \frac{1}{M} \left[\frac{1}{3} \alpha x^3 \right]_{x=0}^{x=L}$$

$$x_{CM} = \frac{1}{M} \left(\frac{1}{3} \alpha L^3 \right) = \frac{1}{M} \left(\frac{2}{3} ML \right) = \frac{2L}{3}$$

Wednesday, Oct. 31, 2007




PHYS 1443-002, Fall 2007
Dr. Jaehoon Yu

11

Center of Mass and Center of Gravity

The center of mass of any symmetric object lies on the axis of symmetry and on any plane of symmetry, if the object's mass is evenly distributed throughout the body.




How do you think you can determine the CM of the objects that are not symmetric?

One can use gravity to locate CM.

1. Hang the object by one point and draw a vertical line following a plum-bob.
2. Hang the object by another point and do the same.
3. The point where the two lines meet is the CM.

Center of Gravity



What does this equation tell you?

Since a rigid object can be considered as a collection of small masses, one can see the total gravitational force exerted on the object as

$$\vec{F}_g = \sum_i \vec{F}_i = \sum_i \Delta m_i \vec{g} = M \vec{g}$$

The net effect of these small gravitational forces is equivalent to a single force acting on a point (**Center of Gravity**) with mass M.

The CoG is the point in an object as if all the gravitational force is acting on!

Motion of a Group of Particles

We've learned that the CM of a system can represent the motion of a system. Therefore, for an isolated system of many particles in which the total mass M is preserved, the velocity, total momentum, acceleration of the system are

Velocity of the system

$$\vec{v}_{CM} = \frac{d\vec{r}_{CM}}{dt} = \frac{d}{dt} \left(\frac{1}{M} \sum m_i \vec{r}_i \right) = \frac{1}{M} \sum m_i \frac{d\vec{r}_i}{dt} = \frac{\sum m_i \vec{v}_i}{M}$$

Total Momentum of the system

$$\vec{p}_{CM} = M \vec{v}_{CM} = M \frac{\sum m_i \vec{v}_i}{M} = \sum m_i \vec{v}_i = \sum \vec{p} = \vec{p}_{tot}$$

Acceleration of the system

$$\vec{a}_{CM} = \frac{d\vec{v}_{CM}}{dt} = \frac{d}{dt} \left(\frac{1}{M} \sum m_i \vec{v}_i \right) = \frac{1}{M} \sum m_i \frac{d\vec{v}_i}{dt} = \frac{\sum m_i \vec{a}_i}{M}$$

The external force exerting on the system

$$\sum \vec{F}_{ext} = M \vec{a}_{CM} = \sum m_i \vec{a}_i = \frac{d\vec{p}_{tot}}{dt}$$

What about the internal forces?

If net external force is 0

$$\sum \vec{F}_{ext} = 0 = \frac{d\vec{p}_{tot}}{dt} \quad \vec{p}_{tot} = \text{const}$$

System's momentum is conserved.