

PHYS 1443 – Section 002

Lecture #25

Wednesday, Dec. 5, 2007

Dr. Jae Yu

- Simple Harmonic Motion
- Equation of the SHM
- Simple Block Spring System
- Energy of the SHO



Announcements

- Final exam
 - Date and time: 11am – 12:30 pm, Monday, Dec. 10
 - Location: SH103
 - Covers: CH9.1 – CH14.3




Vibration or Oscillation

What are the things that vibrate/oscillate?

- Tuning fork
- A pendulum
- A car going over a bump
- Buildings and bridges
- The spider web with a prey

So what is a vibration or oscillation? A periodic motion that repeats over the same path.

A simplest case is a block attached at the end of a coil spring.



Simple Harmonic Motion

Motion that occurs by the force that depends on displacement, and the force is always directed toward the system's equilibrium position.

What is a system that has such characteristics?

A system consists of a mass and a spring

When a spring is stretched from its equilibrium position by a length x , the force acting on the mass is

$$F = -kx$$

It's negative, because the force resists against the change of length, directed toward the equilibrium position.

From Newton's second law

$$F = ma = -kx$$

we obtain

$$a = -\frac{k}{m}x$$

This is a second order differential equation that can be solved but it is beyond the scope of this class.

$$\frac{d^2x}{dt^2} = -\frac{k}{m}x$$

Condition for simple harmonic motion

What do you observe from this equation?

Acceleration is proportional to displacement from the equilibrium.
Acceleration is opposite direction to displacement.

Wednesday, Dec. 5, 2007

This system is doing a simple harmonic motion (SHM).

Dr. Jaehoon Yu

Equation of Simple Harmonic Motion

The solution for the 2nd order differential equation

$$\frac{d^2 x}{dt^2} = -\frac{k}{m} x$$

$$x = A \cos(\omega t + \phi)$$

Generalized expression of a simple harmonic motion

Amplitude

Phase

Angular Frequency

Phase constant

Let's think about the meaning of this equation of motion

What happens when $t=0$ and $\phi=0$?

$$x = A \cos(0 + 0) = A$$

What is ϕ if x is not A at $t=0$?

$$x = A \cos(\phi) = x'$$

$$\phi = \cos^{-1}\left(\frac{x'}{A}\right)$$

An oscillation is fully characterized by its:


What are the maximum/minimum possible values of x ?

$$A/-A$$

- Amplitude
- Period or frequency
- Phase constant



Vibration or Oscillation Properties




The maximum displacement from the equilibrium is

Amplitude

One cycle of the oscillation

The complete to-and-fro motion from an initial point



Period of the motion, T


The time it takes to complete one full cycle

Unit? sec

Frequency of the motion, f

The number of complete cycles per second

Unit? s^{-1}



Relationship between period and frequency?

$$f = \frac{1}{T} \quad \text{or} \quad T = \frac{1}{f}$$

Wednesday, Dec. 5, 2007



PHYS 1443-002, Fall 2007
Dr. Jaehoon Yu

More on Equation of Simple Harmonic Motion

What is the time for full cycle of oscillation?

Since after a full cycle the position must be the same

$$x = A \cos(\omega(t + T) + \phi) = A \cos(\omega t + 2\pi + \phi)$$

The period

$$T = \frac{2\pi}{\omega}$$

One of the properties of an oscillatory motion

How many full cycles of oscillation does this undergo per unit time?

$$f = \frac{1}{T} = \frac{\omega}{2\pi}$$

Frequency

What is the unit?

1/s=Hz

Let's now think about the object's speed and acceleration.

$$x = A \cos(\omega t + \phi)$$

Speed at any given time $v = \frac{dx}{dt} = -\omega A \sin(\omega t + \phi)$ Max speed $v_{\max} = \omega A$

Acceleration at any given time $a = \frac{dv}{dt} = -\omega^2 A \cos(\omega t + \phi) = -\omega^2 x$ Max acceleration $a_{\max} = \omega^2 A$

What do we learn about acceleration?

Acceleration is reverse direction to displacement

Acceleration and speed are $\pi/2$ off phase of each other:

When v is maximum, a is at its minimum



Simple Harmonic Motion continued

Phase constant determines the starting position of a simple harmonic motion.

$$X = A \cos(\omega t + \phi) \quad \boxed{\text{At } t=0} \quad x|_{t=0} = A \cos \phi$$

This constant is important when there are more than one harmonic oscillation involved in the motion and to determine the overall effect of the composite motion

Let's determine phase constant and amplitude

$\boxed{\text{At } t=0}$

$$x_i = A \cos \phi \quad v_i = -\omega A \sin \phi$$

By taking the ratio, one can obtain the phase constant

$$\phi = \tan^{-1} \left(-\frac{v_i}{\omega x_i} \right)$$

By squaring the two equations and adding them together, one can obtain the amplitude

$$x_i^2 = A^2 \cos^2 \phi$$

$$v_i^2 = \omega^2 A^2 \sin^2 \phi$$


$$A^2 (\cos^2 \phi + \sin^2 \phi) = A^2 = x_i^2 + \left(\frac{v_i}{\omega} \right)^2$$

$$A = \sqrt{x_i^2 + \left(\frac{v_i}{\omega} \right)^2}$$



Sinusoidal Behavior of SHM

What do you think the trajectory will look if the oscillation was plotted against time?





Wednesday, Dec. 5, 2007



PHYS 1443-002, Fall 2007
Dr. Jaehoon Yu

Sinusoidal Behavior of SHM



Example for Simple Harmonic Motion

An object oscillates with simple harmonic motion along the x-axis. Its displacement from the origin varies with time according to the equation; $x = (4.00\text{m})\cos\left(\pi t + \frac{\pi}{4}\right)$ where t is in seconds and the angles in the parentheses are in radians. a) Determine the amplitude, frequency, and period of the motion.

From the equation of motion: $X = A\cos(\omega t + \phi) = (4.00\text{m})\cos\left(\pi t + \frac{\pi}{4}\right)$

The amplitude, A, is $A = 4.00\text{m}$ The angular frequency, ω , is $\omega = \pi$

Therefore, frequency and period are $T = \frac{2\pi}{\omega} = \frac{2\pi}{\pi} = 2\text{s}$ $f = \frac{1}{T} = \frac{\omega}{2\pi} = \frac{\pi}{2\pi} = \frac{1}{2}\text{s}^{-1}$

b) Calculate the velocity and acceleration of the object at any time t.

Taking the first derivative on the equation of motion, the velocity is

$$v = \frac{dx}{dt} = -(4.00 \times \pi) \sin\left(\pi t + \frac{\pi}{4}\right) \text{m/s}$$

By the same token, taking the second derivative of equation of motion, the acceleration, a, is

$$a = \frac{d^2x}{dt^2} = -(4.00 \times \pi^2) \cos\left(\pi t + \frac{\pi}{4}\right) \text{m/s}^2$$



Simple Block-Spring System

A block attached at the end of a spring on a frictionless surface experiences acceleration when the spring is displaced from an equilibrium position.

$$\begin{aligned} F_{\text{spring}} &= ma \\ &= -kx \\ a &= -\frac{k}{m}x \end{aligned}$$

This becomes a second order differential equation

$$\frac{d^2x}{dt^2} = -\frac{k}{m}x$$

If we denote

$$\omega^2 = \frac{k}{m}$$

The resulting differential equation becomes

$$\frac{d^2x}{dt^2} = -\omega^2x$$

Since this satisfies condition for simple harmonic motion, we can take the solution

$$x = A \cos(\omega t + \phi)$$

Does this solution satisfy the differential equation?

Let's take derivatives with respect to time $\frac{dx}{dt} = A \frac{d}{dt}(\cos(\omega t + \phi)) = -\omega A \sin(\omega t + \phi)$

Now the second order derivative becomes

$$\frac{d^2x}{dt^2} = -\omega A \frac{d}{dt}(\sin(\omega t + \phi)) = -\omega^2 A \cos(\omega t + \phi) = -\omega^2 x$$

Whenever the force acting on an object is linearly proportional to the displacement from some equilibrium position and is in the opposite direction, the particle moves in simple harmonic motion.



More Simple Block-Spring System

How do the period and frequency of this harmonic motion look?

Since the angular frequency ω is $\omega = \sqrt{\frac{k}{m}}$

The period, T , becomes $T = \frac{2\pi}{\omega} = 2\pi \sqrt{\frac{m}{k}}$

So the frequency is $f = \frac{1}{T} = \frac{\omega}{2\pi} = \frac{1}{2\pi} \sqrt{\frac{k}{m}}$

What can we learn from these?

- Frequency and period do not depend on amplitude
- Period is inversely proportional to spring constant and proportional to mass

Special case #1

Let's consider that the spring is stretched to a distance A , and the block is let go from rest, giving 0 initial speed; $x_i = A$, $v_i = 0$,

$$x = A \cos \omega t \quad v = \frac{dx}{dt} = -\omega A \sin \omega t \quad a = \frac{d^2x}{dt^2} = -\omega^2 A \cos \omega t \quad a_i = -\omega^2 A = -kA/m$$

This equation of motion satisfies all the conditions. So it is the solution for this motion.

Special case #2

Suppose a block is given non-zero initial velocity v_i to positive x at the instant it is at the equilibrium, $x_i = 0$

$$\phi = \tan^{-1}\left(-\frac{v_i}{\omega x_i}\right) = \tan^{-1}(-\infty) = -\frac{\pi}{2} \quad x = A \cos\left(\omega t - \frac{\pi}{2}\right) = A \sin(\omega t)$$

Is this a good solution?



Example for Spring Block System

A car with a mass of 1300kg is constructed so that its frame is supported by four springs. Each spring has a force constant of 20,000N/m. If two people riding in the car have a combined mass of 160kg, find the frequency of vibration of the car after it is driven over a pothole in the road.

Let's assume that mass is evenly distributed to all four springs.

The total mass of the system is 1460kg.

Therefore each spring supports 365kg each.

From the frequency relationship
based on Hook's law

$$f = \frac{1}{T} = \frac{\omega}{2\pi} = \frac{1}{2\pi} \sqrt{\frac{k}{m}}$$

Thus the frequency for
vibration of each spring is

$$f = \frac{1}{2\pi} \sqrt{\frac{k}{m}} = \frac{1}{2\pi} \sqrt{\frac{20000}{365}} = 1.18 \text{ s}^{-1} = 1.18 \text{ Hz}$$


How long does it take for the car to complete two full vibrations?

The period is $T = \frac{1}{f} = 2\pi \sqrt{\frac{m}{k}} = 0.849 \text{ s}$ For two cycles $2T = 1.70 \text{ s}$



Example for Spring Block System

A block with a mass of 200g is connected to a light spring for which the force constant is 5.00 N/m and is free to oscillate on a horizontal, frictionless surface. The block is displaced 5.00 cm from equilibrium and released from rest. Find the period of its motion.



From the Hook's law, we obtain

$$\omega = \sqrt{\frac{k}{m}} = \sqrt{\frac{5.00}{0.20}} = 5.00 \text{ s}^{-1}$$

As we know, period does not depend on the amplitude or phase constant of the oscillation, therefore the period, T , is simply

$$T = \frac{2\pi}{\omega} = \frac{2\pi}{5.00} = 1.26 \text{ s}$$

Determine the maximum speed of the block.

From the general expression of the simple harmonic motion, the speed is

$$\begin{aligned} v_{\max} &= \frac{dx}{dt} = -\omega A \sin(\omega t + \phi) \\ &= \omega A = 5.00 \times 0.05 = 0.25 \text{ m/s} \end{aligned}$$

Energy of the Simple Harmonic Oscillator

What do you think the mechanical energy of the harmonic oscillator look without friction?


Kinetic energy of a harmonic oscillator is

$$KE = \frac{1}{2}mv^2 = \frac{1}{2}m\omega^2 A^2 \sin^2(\omega t + \phi)$$

The elastic potential energy stored in the spring $PE = \frac{1}{2}kx^2 = \frac{1}{2}kA^2 \cos^2(\omega t + \phi)$

Therefore the total mechanical energy of the harmonic oscillator is $E = KE + PE = \frac{1}{2}[m\omega^2 A^2 \sin^2(\omega t + \phi) + kA^2 \cos^2(\omega t + \phi)]$

Since $\omega = \sqrt{k/m}$ $E = KE + PE = \frac{1}{2}[kA^2 \sin^2(\omega t + \phi) + kA^2 \cos^2(\omega t + \phi)] = \frac{1}{2}kA^2$



Total mechanical energy of a simple harmonic oscillator is proportional to the square of the amplitude.

Energy of the Simple Harmonic Oscillator cont'd

Maximum KE is
when PE=0

$$KE_{\max} = \frac{1}{2} m v_{\max}^2 = \frac{1}{2} k A^2$$


Maximum speed

$$v_{\max} = \sqrt{\frac{k}{m}} A$$


*The speed at any given
point of the oscillation*

$$E = KE + PE = \frac{1}{2} m v^2 + \frac{1}{2} k x^2 = \frac{1}{2} k A^2$$

$$v = \sqrt{k/m (A^2 - x^2)} = v_{\max} \sqrt{1 - \left(\frac{x}{A}\right)^2}$$



Oscillation Properties



Amplitude? **A**

- When is the force greatest?
- When is the speed greatest?
- When is the acceleration greatest?
- When is the potential energy greatest?
- When is the kinetic energy greatest?

Example for Energy of Simple Harmonic Oscillator

A 0.500kg cube connected to a light spring for which the force constant is 20.0 N/m oscillates on a horizontal, frictionless track. a) Calculate the total energy of the system and the maximum speed of the cube if the amplitude of the motion is 3.00 cm.

From the problem statement, A and k are

$$k = 20.0 \text{ N / m}$$

$$A = 3.00 \text{ cm} = 0.03 \text{ m}$$

The total energy of the cube is

$$E = KE + PE = \frac{1}{2} k A^2 = \frac{1}{2} (20.0) \times (0.03)^2 = 9.00 \times 10^{-3} \text{ J}$$

Maximum speed occurs when kinetic energy is the same as the total energy

$$KE_{\max} = \frac{1}{2} m v_{\max}^2 = E = \frac{1}{2} k A^2$$

$$v_{\max} = A \sqrt{\frac{k}{m}} = 0.03 \sqrt{\frac{20.0}{0.500}} = 0.190 \text{ m / s}$$

b) What is the velocity of the cube when the displacement is 2.00 cm.

velocity at any given displacement is

$$V = \sqrt{k/m(A^2 - x^2)} = \sqrt{20.0 \cdot (0.03^2 - 0.02^2) / 0.500} = 0.141 \text{ m / s}$$

c) Compute the kinetic and potential energies of the system when the displacement is 2.00 cm.

Kinetic energy, KE $KE = \frac{1}{2} m v^2 = \frac{1}{2} 0.500 \times (0.141)^2 = 4.97 \times 10^{-3} \text{ J}$ Potential energy, PE $PE = \frac{1}{2} k x^2 = \frac{1}{2} 20.0 \times (0.02)^2 = 4.00 \times 10^{-3} \text{ J}$



Congratulations!!!!

You all have done very well!!!

*I certainly had a lot of fun with ya'll
and am truly proud of you!*

Good luck with your exam!!!

Have safe holidays!!

