

PHYS 1443 – Section 002

Lecture #19

Monday, Nov. 24, 2008

Dr. Jae Yu

- Rolling Motion of a Rigid Body
- Angular Momentum
- Conservation of Angular Momentum
- Relationship between angular and linear quantities
- Similarity between Linear and Angular Quantities

Today's homework is HW #11, due 9pm, Monday, Dec. 1!!



Announcements

- 3rd term exam grading not completed
 - Will be done by Monday, Dec. 1
- Thanksgiving is this Thursday, Nov. 27
 - Vote for the class this Wednesday, Nov. 26



Rolling Motion of a Rigid Body

What is a rolling motion?

A more generalized case of a motion where the rotational axis moves together with an object

A rotational motion about a moving axis


To simplify the discussion, let's make a few assumptions

1. Limit our discussion on very symmetric objects, such as cylinders, spheres, etc
2. The object rolls on a flat surface

Let's consider a cylinder rolling on a flat surface, without slipping.

Under what condition does this "Pure Rolling" happen?

The total linear distance the CM of the cylinder moved is $s = R\theta$



Thus the linear speed of the CM is


$$v_{CM} = \frac{ds}{dt} = R \frac{d\theta}{dt} = R\omega$$

The condition for a "Pure Rolling motion"

More Rolling Motion of a Rigid Body

The magnitude of the linear acceleration of the CM is

$$a_{CM} = \frac{dv_{CM}}{dt} = R \frac{d\omega}{dt} = R\alpha$$




As we learned in rotational motion, all points in a rigid body moves at the same angular speed but at different linear speeds.

CM is moving at the same speed at all times.


At any given time, the point that comes to P has 0 linear speed while the point at P' has twice the speed of CM

Why??

A rolling motion can be interpreted as the sum of Translation and Rotation



Kinetic Energy of a Rolling Sphere



Let's consider a sphere with radius R rolling down the hill without slipping.

Since $v_{CM} = R\omega$

$$\begin{aligned} K &= \frac{1}{2} I_{CM} \omega^2 + \frac{1}{2} M R^2 \omega^2 \\ &= \frac{1}{2} I_{CM} \left(\frac{v_{CM}}{R} \right)^2 + \frac{1}{2} M v_{CM}^2 \\ &= \frac{1}{2} \left(\frac{I_{CM}}{R^2} + M \right) v_{CM}^2 \end{aligned}$$

What is the speed of the CM in terms of known quantities and how do you find this out?


Since the kinetic energy at the bottom of the hill must be equal to the potential energy at the top of the hill

$$K = \frac{1}{2} \left(\frac{I_{CM}}{R^2} + M \right) v_{CM}^2 = Mgh$$

$$v_{CM} = \sqrt{\frac{2gh}{1 + I_{CM} / MR^2}}$$

Example for Rolling Kinetic Energy

For solid sphere as shown in the figure, calculate the linear speed of the CM at the bottom of the hill and the magnitude of linear acceleration of the CM. Solve this problem using Newton's second law, the dynamic method.



What are the forces involved in this motion?

Gravitational Force, Frictional Force, Normal Force

Newton's second law applied to the CM gives

$$\begin{aligned}\sum F_x &= Mg \sin \theta - f = Ma_{CM} \\ \sum F_y &= n - Mg \cos \theta = 0\end{aligned}$$

Since the forces Mg and n go through the CM, their moment arm is 0 and do not contribute to torque, while the static friction f causes torque $\tau_{CM} = fR = I_{CM}\alpha$

We know that

$$I_{CM} = \frac{2}{5}MR^2$$

$$a_{CM} = R\alpha$$

We obtain


Substituting f in dynamic equations

$$f = \frac{I_{CM}\alpha}{R} = \frac{\frac{2}{5}MR^2}{R} \left(\frac{a_{CM}}{R} \right) = \frac{2}{5}Ma_{CM}$$

$$Mg \sin \theta = \frac{7}{5}Ma_{CM} \quad a_{CM} = \frac{5}{7}g \sin \theta$$

Angular Momentum of a Particle

If you grab onto a pole while running, your body will rotate about the pole, gaining angular momentum. We've used the linear momentum to solve physical problems with linear motions, the angular momentum will do the same for rotational motions.



Let's consider a point-like object (particle) with mass m located at the vector location \mathbf{r} and moving with linear velocity \mathbf{v}

The angular momentum \mathcal{L} of this particle relative to the origin O is

$$\vec{\mathbf{L}} \equiv \vec{\mathbf{r}} \times \vec{\mathbf{p}}$$

What is the unit and dimension of angular momentum? $\text{kg} \cdot \text{m}^2 / \text{s}$ $[\text{ML}^2\text{T}^{-1}]$

Note that \mathcal{L} depends on origin O. Why? Because \mathbf{r} changes

What else do you learn? The direction of \mathcal{L} is +z.

Since \mathbf{p} is $m\mathbf{v}$, the magnitude of \mathcal{L} becomes $L = mvr \sin \phi$

What do you learn from this?

If the direction of linear velocity points to the origin of rotation, the particle does not have any angular momentum.

If the linear velocity is perpendicular to position vector, the particle moves exactly the same way as a point on a rim.

Angular Momentum and Torque


Can you remember how net force exerting on a particle and the change of its linear momentum are related?

$$\sum \vec{F} = \frac{d\vec{p}}{dt}$$

Total external forces exerting on a particle is the same as the change of its linear momentum.

The same analogy works in rotational motion between torque and angular momentum.

Net torque acting on the particle is $\sum \vec{\tau} = \vec{r} \times \sum \vec{F} = \vec{r} \times \frac{d\vec{p}}{dt}$



$$\frac{d\vec{L}}{dt} = \frac{d(\vec{r} \times \vec{p})}{dt} = \frac{d\vec{r}}{dt} \times \vec{p} + \vec{r} \times \frac{d\vec{p}}{dt} = 0 + \vec{r} \times \frac{d\vec{p}}{dt} = \sum \vec{\tau}$$

Why does this work?

Because \vec{v} is parallel to the linear momentum

Thus the torque-angular momentum relationship

$$\sum \vec{\tau} = \frac{d\vec{L}}{dt}$$

The net torque acting on a particle is the same as the time rate change of its angular momentum

Monday, Nov. 24, 2008



PHYS 1443-002, Fall 2006 Dr. Jaehoon Yu

Yu

o

Angular Momentum of a System of Particles

The total angular momentum of a system of particles about some point is the vector sum of the angular momenta of the individual particles

$$\vec{L} = \vec{L}_1 + \vec{L}_2 + \dots + \vec{L}_n = \sum \vec{L}_i$$

Since the individual angular momentum can change, the total angular momentum of the system can change.

Both internal and external forces can provide torque to individual particles. However, the internal forces do not generate net torque due to Newton's third law.

Let's consider a two particle system where the two exert forces on each other.

Since these forces are the action and reaction forces with directions lie on the line connecting the two particles, the vector sum of the torque from these two becomes 0.


Thus the time rate change of the angular momentum of a system of particles is equal to only the net external torque acting on the system

$$\sum \vec{\tau}_{ext} = \frac{d\vec{L}}{dt}$$



Example for Angular Momentum

A particle of mass m is moving on the xy plane in a circular path of radius r and linear velocity v about the origin O . Find the magnitude and the direction of the angular momentum with respect to O .



Using the definition of angular momentum

$$\vec{L} = \vec{r} \times \vec{p} = \vec{r} \times m \vec{v} = m \vec{r} \times \vec{v}$$

Since both the vectors, \vec{r} and \vec{v} , are on x - y plane and using right-hand rule, the direction of the angular momentum vector is $+z$ (coming out of the screen)

The magnitude of the angular momentum is $|\vec{L}| = |\vec{r} \times m\vec{v}| = mrv \sin \phi = mrv \sin 90^\circ = mrv$


So the angular momentum vector can be expressed as $\vec{L} = mrv \vec{k}$

Find the angular momentum in terms of angular velocity ω .

Using the relationship between linear and angular speed

$$\vec{L} = mrv \vec{k} = mr^2 \omega \vec{k} = mr^2 \vec{\omega} = I \vec{\omega}$$

Angular Momentum of a Rotating Rigid Body



Let's consider a rigid body rotating about a fixed axis

Each particle of the object rotates in the xy plane about the z-axis at the same angular speed, ω

Magnitude of the angular momentum of a particle of mass m_i about origin O is $m_i v_i r_i$ $L_i = m_i r_i v_i = m_i r_i^2 \omega$

Summing over all particle's angular momentum about z axis

$$L_z = \sum_i L_i = \sum_i (m_i r_i^2 \omega)$$

What do you see?

$$L_z = \sum_i (m_i r_i^2) \omega = I \omega$$

Since I is constant for a rigid body

$$\frac{dL_z}{dt} = I \frac{d\omega}{dt} = I \alpha$$

α is angular acceleration


Thus the torque-angular momentum relationship becomes

$$\sum \tau_{ext} = \frac{dL_z}{dt} = I \alpha$$

The net external torque acting on a rigid body rotating about a fixed axis is equal to the moment of inertia about that axis multiplied by the object's angular acceleration with respect to that axis.

Example for Rigid Body Angular Momentum

A rigid rod of mass \mathcal{M} and length ℓ is pivoted without friction at its center. Two particles of mass m_1 and m_2 are attached to either end of the rod. The combination rotates on a vertical plane with an angular speed of ω . Find an expression for the magnitude of the angular momentum.



The moment of inertia of this system is

$$I = I_{rod} + I_{m_1} + I_{m_2} = \frac{1}{12} M l^2 + \frac{1}{4} m_1 l^2 + \frac{1}{4} m_2 l^2$$

$$= \frac{l^2}{4} \left(\frac{1}{3} M + m_1 + m_2 \right)$$

$$L = I \omega = \frac{\omega l^2}{4} \left(\frac{1}{3} M + m_1 + m_2 \right)$$

Find an expression for the magnitude of the angular acceleration of the system when the rod makes an angle θ with the horizon.

If $m_1 = m_2$, no angular momentum because the net torque is 0.

If $\theta = \pm \pi/2$, at equilibrium so no angular momentum.

First compute the net external torque

$$\tau_1 = m_1 g \frac{l}{2} \cos \theta \quad \tau_2 = -m_2 g \frac{l}{2} \cos \theta$$

$$\tau_{ext} = \tau_1 + \tau_2 = \frac{gl \cos \theta (m_1 - m_2)}{2}$$

Thus α becomes

$$\alpha = \frac{\sum \tau_{ext}}{I} = \frac{\frac{1}{2} (m_1 - m_2) gl \cos \theta}{\frac{l^2}{4} \left(\frac{1}{3} M + m_1 + m_2 \right)} = \frac{2 (m_1 - m_2) \cos \theta}{\left(\frac{1}{3} M + m_1 + m_2 \right)} \frac{g}{l}$$

Monday, Nov. 24, 2008



-002, Fall 2008 Dr. Yu

Conservation of Angular Momentum

Remember under what condition the linear momentum is conserved?

Linear momentum is conserved when the net external force is 0. $\sum \vec{F} = 0 = \frac{d\vec{p}}{dt}$
 $\vec{p} = \text{const}$

By the same token, the angular momentum of a system is constant in both magnitude and direction, if the resultant external torque acting on the system is 0.

$$\sum \vec{\tau}_{ext} = \frac{d\vec{L}}{dt} = 0$$

$$\vec{L} = \text{const}$$

What does this mean?

Angular momentum of the system before and after a certain change is the same.

$$\vec{L}_i = \vec{L}_f = \text{constant}$$

Three important conservation laws for isolated system that does not get affected by external forces

$$K_i + U_i = K_f + U_f$$

$$\vec{p}_i = \vec{p}_f$$

$$\vec{L}_i = \vec{L}_f$$

Mechanical Energy

Linear Momentum

Angular Momentum



Example for Angular Momentum Conservation

A star rotates with a period of 30 days about an axis through its center. After the star undergoes a supernova explosion, the stellar core, which had a radius of $1.0 \times 10^4 \text{ km}$, collapses into a neutron star of radius 3.0 km . Determine the period of rotation of the neutron star.

What is your guess about the answer?

The period will be significantly shorter, because its radius got smaller.

Let's make some assumptions:

1. There is no external torque acting on it
2. The shape remains spherical
3. Its mass remains constant

Using angular momentum conservation

$$L_i = L_f$$

$$I_i \omega_i = I_f \omega_f$$

The angular speed of the star with the period T is

$$\omega = \frac{2\pi}{T}$$


Thus
$$\omega_f = \frac{I_i \omega_i}{I_f} = \frac{mr_i^2}{mr_f^2} \frac{2\pi}{T_i}$$

$$T_f = \frac{2\pi}{\omega_f} = \left(\frac{r_f^2}{r_i^2} \right) T_i = \left(\frac{3.0}{1.0 \times 10^4} \right)^2 \times 30 \text{ days} = 2.7 \times 10^{-6} \text{ days} = 0.23 \text{ s}$$



Kepler's Second Law and Angular Momentum Conservation

Consider a planet of mass M_p moving around the Sun in an elliptical orbit.



Since the gravitational force acting on the planet is always toward radial direction, it is a *central force*. Therefore the torque acting on the planet by this force is always 0.

$$\vec{\tau} = \vec{r} \times \vec{F} = \vec{r} \times F\hat{r} = 0$$

Since torque is the time rate change of angular momentum \vec{L} , the angular momentum is constant.

$$\vec{\tau} = \frac{d\vec{L}}{dt} = 0 \quad \vec{L} = \text{const}$$

Because the gravitational force exerted on a planet by the Sun results in no torque, the angular momentum L of the planet is constant.

$$\vec{L} = \vec{r} \times \vec{p} = \vec{r} \times M_p \vec{v} = M_p \vec{r} \times \vec{v} = \text{const}$$

Since the area swept by the motion of the planet is

$$dA = \frac{1}{2} |\vec{r} \times d\vec{r}| = \frac{1}{2} |\vec{r} \times \vec{v} dt| = \frac{L}{2M_p} dt \quad \Rightarrow \quad \frac{dA}{dt} = \frac{L}{2M_p} = \text{const}$$

This is Kepler's second law which states that the radius vector from the Sun to a planet sweeps out equal areas in equal time intervals.

Similarity Between Linear and Rotational Motions

All physical quantities in linear and rotational motions show striking similarity.

Quantities	Linear	Rotational
Mass	Mass M	Moment of Inertia $I = mr^2$
Length of motion	Distance L	Angle θ (Radian)
Speed	$v = \frac{\Delta r}{\Delta t}$	$\omega = \frac{\Delta \theta}{\Delta t}$
Acceleration	$a = \frac{\Delta v}{\Delta t}$	$\alpha = \frac{\Delta \omega}{\Delta t}$
Force	Force $\vec{F} = m\vec{a}$	Torque $\vec{\tau} = I\vec{\alpha}$
Work	Work $W = \vec{F} \cdot \vec{d}$	Work $W = \tau\theta$
Power	$P = \vec{F} \cdot \vec{v}$	$P = \tau\omega$
Momentum	$\vec{p} = m\vec{v}$	$\vec{L} = I\vec{\omega}$
Kinetic Energy	Kinetic $K = \frac{1}{2}mv^2$	Rotational $K_R = \frac{1}{2}I\omega^2$

