

PHYS 1444 – Section 002

Lecture #6

Monday, Feb. 10, 2020

Dr. Jaehoon Yu

- Ch 21
 - Motion of a Charged Particle in an E Field
 - Electric Dipole and Dipole Moment
- CH 22
 - Gauss' Law
 - Electric Flux
 - Gauss' Law with Multiple Charges

Today's homework is homework #4, due 11pm, Monday, Feb. 17!!



Announcements – I

- Please make sure you make the necessary payment for the homework
 - You and I both will lose access to your homework!
- Quiz # 2
 - Beginning of the class this Wednesday, Feb. 12
 - Covers CH21.6 to what we finish today (CH22?)
 - You can bring your calculator w/o any relevant formula pre-input
 - BYOF: You may bring a one 8.5x11.5 sheet (front and back) of handwritten formulae and values of constants for the exam
 - No derivations, word definitions, or solutions of any problems !
 - No additional formulae or values of constants will be provided!



Announcements – II

- 1st Term Exam
 - In class, Wednesday, Feb. 19: DO NOT MISS THE EXAM!
 - CH21.1 to what we learn on Monday, Feb. 17 + Appendices A1 – A8
 - You can bring your calculator w/o any relevant formula pre-input
 - BYOF: You may bring a one 8.5x11.5 sheet (front and back) of handwritten formulae and values of constants for the exam
 - No derivations, word definitions, or solutions of any problems !
 - No additional formulae or values of constants will be provided!
- Quiz 1 results
 - Class average: 38.7/80
 - Equivalent to 48.4/100
 - Top score: 73/80



Reminder: Special Project #2 – Angels & Demons

- Compute the total possible energy released from an annihilation of xx-grams of anti-matter and the same quantity of matter, where xx is the last two digits of your SS#. (20 points)
 - Use the famous Einstein's formula for mass-energy equivalence
- Compute the power output of this annihilation when the energy is released in yy ns, where yy is the first two digits of your SS#. (10 points)
- Compute how many cups of gasoline (8MJ) this energy corresponds to. (5 points)
- Compute how many months of world electricity usage (3.6GJ/mo) this energy corresponds to. (5 points)
- Due at the beginning of the class Monday, Feb. 24


Monday, Feb. 10, 2020



PHYS 1444-002, Spring 2020
Dr. Jaehoon Yu

What did we last weekend?



- Built the first field cage module for DUNE!



Monday, Feb. 10, 2020


Motion of a Charged Particle in an Electric Field

- If an object with an electric charge q is at a point in space where electric field is \mathbf{E} , the force exerting on the object is $\vec{F} = q\vec{E}$.
- What do you think will happen to the charge?
 - Let's think about the cases like these on the right.
 - The object will move along the field line...Which way?
 - Depends on the sign of the charge
 - The charge gets accelerated under an electric field.



Example 21 – 14

- Electron accelerated by electric field.** An electron (mass $m = 9.1 \times 10^{-31} \text{ kg}$) is accelerated in a uniform field E ($E = 2.0 \times 10^4 \text{ N/C}$) between two parallel charged plates. The separation of the plates is 1.5 cm. The electron is accelerated from rest near the negative plate and passes through a tiny hole in the positive plate. (a) With what speed does it leave the hole? (b) Show that the gravitational force can be ignored. Assume the hole is so small that it does not affect the uniform field between the plates.



The magnitude of the force on the electron is $F = qE$ and is directed to the right. The equation to solve this problem is

$$F = qE = ma$$

The magnitude of the electron's acceleration is $a = \frac{F}{m} = \frac{qE}{m}$

Between the plates the field E is uniform, thus the electron undergoes a uniform acceleration

$$a = \frac{eE}{m_e} = \frac{(1.6 \times 10^{-19} \text{ C})(2.0 \times 10^4 \text{ N/C})}{(9.1 \times 10^{-31} \text{ kg})} = 3.5 \times 10^{15} \text{ m/s}^2$$

Example 21 – 14

Since the travel distance is $1.5 \times 10^{-2} \text{m}$, using one of the kinetic eq. of motions,

$$v^2 = v_0^2 + 2ax \quad \therefore v = \sqrt{2ax} = \sqrt{2 \cdot 3.5 \times 10^{15} \cdot 1.5 \times 10^{-2}} = 1.0 \times 10^7 \text{ m/s}$$

Since there is no electric field outside the conductor, the electron continues moving with this speed after passing through the hole.

- (b) Show that the gravitational force can be ignored. Assume the hole is so small that it does not affect the uniform field between the plates.

The magnitude of the electric force on the electron is

$$F_e = qE = eE = (1.6 \times 10^{-19} \text{ C})(2.0 \times 10^4 \text{ N/C}) = 3.2 \times 10^{-15} \text{ N}$$


The magnitude of the gravitational force on the electron is

$$F_G = mg = 9.8 \text{ m/s}^2 \times (9.1 \times 10^{-31} \text{ kg}) = 8.9 \times 10^{-30} \text{ N}$$


Thus the gravitational force on the electron is negligible compared to the electromagnetic force.

Electric Dipoles

- An electric dipole is the combination of two equal charges of opposite signs, $+Q$ and $-Q$, separated by a distance ℓ , which behaves as one entity.
- The quantity $Q\ell$ is called the electric dipole moment and is represented by the symbol p .




- The dipole moment is a vector quantity, p
- The magnitude of the dipole moment is $Q\ell$ Unit? **C-m**
- Its direction is from the negative to the positive charge.
- Many of diatomic molecules like CO have a dipole moment. \rightarrow These are referred as polar molecules.
 - Even if the molecule is electrically neutral, their sharing of electron causes separation of charges
 - Symmetric diatomic molecules, such as O_2 , do not have a dipole moment
- The water molecule also has a dipole moment which is the vector sum of two dipole moments between Oxygen and each of Hydrogen atoms.




Dipoles in an External Field

- Let's consider a dipole placed in a uniform electric field \mathbf{E} .
- What do you think will happen to the dipole in the figure?
 - Forces will be exerted on the charges.
 - The positive charge will get pushed toward right while the negative charge will get pulled toward left.
 - What is the net force acting on the dipole?
 - Zero
 - So will the dipole not move?
 - Yes, it will.
 - Why?
 - There is a torque applied on the dipole.



Dipoles in an External Field, cnt'd

- How much is the torque on the dipole?
 - Do you remember the formula for torque?
 - $\vec{\tau} = \vec{r} \times \vec{F}$
 - The magnitude of the torque exerting on each of the charges with respect to the rotational axis at the center is
 - $\tau_{+Q} = |\vec{r} \times \vec{F}| = rF \sin \theta = \left(\frac{l}{2} \right) (QE) \sin \theta = \frac{l}{2} QE \sin \theta$
 - $\tau_{-Q} = |\vec{r} \times \vec{F}| = rF \sin \theta = \left(-\frac{l}{2} \right) (-QE) \sin \theta = \frac{l}{2} QE \sin \theta$
 - Thus, the total torque is
 - $\tau_{Total} = \tau_{+Q} + \tau_{-Q} = \frac{l}{2} QE \sin \theta + \frac{l}{2} QE \sin \theta = lQE \sin \theta = pE \sin \theta$
 - So the torque on a dipole in vector notation is $\vec{\tau} = \vec{p} \times \vec{E}$
- The effect of the torque is to try to turn the dipole so that the dipole moment is parallel to \vec{E} . Which direction?



Potential Energy of a Dipole in an External Field

- What is the work done on the dipole by the electric field to change the angle from θ_1 to θ_2 ?

$$W = \int_{\theta_1}^{\theta_2} dW = \int_{\theta_1}^{\theta_2} \vec{\tau} \cdot d\vec{\theta} = \int_{\theta_1}^{\theta_2} -\tau d\theta$$

Why negative?

Because τ and θ are in opposite directions to each other.

- The torque is $\tau = pE \sin \theta$.

- Thus the work done on the dipole by the field is

$$W = \int_{\theta_1}^{\theta_2} -pE \sin \theta d\theta = pE [\cos \theta]_{\theta_1}^{\theta_2} = pE (\cos \theta_2 - \cos \theta_1)$$

- What happens to the dipole's potential energy, U , when a positive work is done on it by the field?

– It decreases.

- We choose $U=0$ when $\theta_1=90$ degrees, then the potential energy at $\theta_2=\theta$ becomes $U = -W = -pE \cos \theta = -\vec{p} \cdot \vec{E}$


Electric Field by a Dipole

- Let's consider the case in the picture.
- There are fields by both the charges. So the total electric field by the dipole is $\vec{E}_{Tot} = \vec{E}_{+Q} + \vec{E}_{-Q}$
- The magnitudes of the two fields are equal

$$E_{+Q} = E_{-Q} = \frac{1}{4\pi\epsilon_0} \frac{Q}{\left(\sqrt{r^2 + (l/2)^2}\right)^2} = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2 + (l/2)^2} = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2 + l^2/4}$$

- Now we must work out the x and y components of the total field.
 - Sum of the two y components is
 - Zero since they are the same but in opposite direction
 - So the magnitude of the total field is the same as the sum of the two x-components:

$$E = 2E_+ \cos \phi = \frac{1}{2\pi\epsilon_0} \frac{Q}{r^2 + l^2/4} \frac{l}{2\sqrt{r^2 + l^2/4}} = \frac{1}{4\pi\epsilon_0} \frac{p}{\left(r^2 + l^2/4\right)^{3/2}}$$



Dipole Electric Field from Afar

- What happens when $r \gg l$?

$$E_D = \frac{1}{4\pi\epsilon_0} \frac{p}{\left(r^2 + l^2/4\right)^{3/2}} \approx \frac{1}{4\pi\epsilon_0} \frac{p}{r^3} \quad (\text{when } r \gg l)$$

- Why does this make sense?
 - Since from a long distance, the two charges are very close so that the overall charge gets close to 0!!
 - This dependence works for the point not on the bisecting line as well



Example 21 – 17

- **Dipole in a field.** The dipole moment of a water molecule is $6.1 \times 10^{-30} \text{C}\cdot\text{m}$. A water molecule is placed in a uniform electric field with magnitude $2.0 \times 10^5 \text{N/C}$. (a) What is the magnitude of the maximum torque that the field can exert on the molecule? (b) What is the potential energy when the torque is at its maximum? (c) In what position will the potential energy take on its greatest value? Why is this different than the position where the torque is maximized?

(a) The torque is maximized when $\theta = 90$ degrees. Thus the magnitude of the maximum torque is

$$\begin{aligned}\tau &= pE \sin \theta = pE = \\ &= (6.1 \times 10^{-30} \text{C} \cdot \text{m})(2.5 \times 10^5 \text{N/C}) = 1.2 \times 10^{-24} \text{N} \cdot \text{m}\end{aligned}$$

What is the distance between a hydrogen atom and the oxygen atom?

Example 21 – 17

(b) What is the potential energy when the torque is at its maximum?

Since the dipole potential energy is $U = -\vec{p} \cdot \vec{E} = -pE \cos \theta$

And τ is at its maximum at $\theta=90$ degrees, the potential energy, U , is

$$U = -pE \cos \theta = -pE \cos(90^\circ) = 0$$

Is the potential energy at its minimum at $\theta=90$ degrees? **No**

Why not? **Because U will become negative as θ increases.**

(c) In what position will the potential energy take on its greatest value?

The potential energy is maximum when $\cos\theta = -1$, $\theta=180$ degrees.

Why is this different than the position where the torque is maximized?

The potential energy is maximized when the dipole is oriented so that it has to rotate through the largest angle against the direction of the field, to reach the equilibrium position at $\theta=0$.

Torque is maximized when the field is perpendicular to the dipole, $\theta=90$.




Gauss' Law

- Gauss' law states the relationship between electric charge and the electric field.
 - More generalized and elegant form of Coulomb's law.
- The electric field by a distribution of charges can be obtained using Coulomb's law by summing (or integrating) over the charge distributions vectorially.
- Gauss' law, however, gives an additional insight into the nature of the electrostatic field and a more general relationship between the charge and the field




Electric Flux



- Let's imagine a surface of area A through which a uniform electric field \mathbf{E} passes
- The electric flux Φ_E is defined as
 - $\Phi_E = EA$, if the field is perpendicular to the surface
 - $\Phi_E = EA \cos \theta$, if the field makes an angle θ to the surface
- So the electric flux is defined as $\Phi_E = \vec{E} \cdot \vec{A}$. **Unit?** $\text{N} \cdot \text{m}^2 / \text{C}$
- How would you define the electric flux in words?
 - The total number of field lines passing through the unit area perpendicular to the field. $N_E \propto EA_{\perp} = \Phi_E$

Example 22 – 1

- Electric flux.** (a) Calculate the electric flux through the rectangle in the figure (a). The rectangle is 10cm by 20cm and the electric field is uniform with magnitude 200N/C. (b) What is the flux in figure if the angle is 30 degrees?



The electric flux is defined as

$$\Phi_E = \vec{E} \cdot \vec{A} = EA \cos \theta$$

So when (a) $\theta=0$, we obtain

$$\Phi_E = EA \cos \theta = EA = (200 \text{ N/C}) \cdot (0.1 \times 0.2 \text{ m}^2) = 4.0 \text{ N} \cdot \text{m}^2/\text{C}$$

And when (b) $\theta=30$ degrees, we obtain

$$\Phi_E = EA \cos 30^\circ = (200 \text{ N/C}) \cdot (0.1 \times 0.2 \text{ m}^2) \cos 30^\circ = 3.5 \text{ N} \cdot \text{m}^2/\text{C}$$



A Brain Teaser of Electric Flux

- What would change the electric flux through a circle lying in the xz plane where the electric field is $(10\text{N/C})\mathbf{j}$?
 1. Changing the magnitude of the electric field
 2. Changing the surface area of the circle
 3. Tipping the circle so that it is lying in the xy plane
 4. All of the above
 5. None of the above



Generalization of the Electric Flux

- Let's consider a surface of area A that is not a square or flat but in some random shape, and that the field is not uniform.
- The surface can be divided up into infinitesimally small areas of $\Delta\mathbf{A}_i$ that can be considered flat.
- And the electric field through this area can be considered uniform since the area is very small.
- Then the electric flux through the entire surface is approximately
- In the limit where $\Delta\mathbf{A}_i \rightarrow 0$, the discrete summation becomes an integral.




$$\Phi_E \approx \sum_{i=1}^n \vec{E}_i \cdot \Delta\vec{A}_i$$

$$\Phi_E = \int \vec{E}_i \cdot d\vec{A} \quad \text{open surface}$$


$$\Phi_E = \oint \vec{E}_i \cdot d\vec{A} \quad \text{enclosed surface}$$

Generalization of the Electric Flux

- We arbitrarily define that the area vector points outward from the enclosed volume.





- For the line leaving the volume, $\theta < \pi/2$ and $\cos\theta > 0$. The flux is positive.
- For the line coming into the volume, $\theta > \pi/2$ and $\cos\theta < 0$. The flux is negative.
- If $\Phi_E > 0$, there is net flux out of the volume.
- If $\Phi_E < 0$, there is flux into the volume.




- In the above figures, each field that enters the volume also leaves the volume, so $\Phi_E = \oint \vec{E} \cdot d\vec{A} = 0$.
- The flux is non-zero only if one or more lines start or end inside the surface.

Generalization of the Electric Flux

- The field line starts or ends only on a charge.
- Sign of the net flux on the surface A_1 ?
 - The net outward flux (positive flux)
- How about A_2 ?
 - Net inward flux (negative flux)
- What is the flux in the bottom figure?
 - There should be a net inward flux (negative flux) since the total charge inside the volume is negative.
- The net flux that crosses an enclosed surface is proportional to the total charge inside the surface. → This is the crux of Gauss' law.



Gauss' Law




- Let's consider the case in the above figure.
- What are the results of the closed integral of the Gaussian surfaces A_1 and A_2 ?

– For A_1 $\oint \vec{E} \cdot d\vec{A} = \frac{+q}{\epsilon_0}$

– For A_2 $\oint \vec{E} \cdot d\vec{A} = \frac{-q'}{\epsilon_0}$

Coulomb's Law from Gauss' Law



- Let's consider a charge Q enclosed inside our imaginary Gaussian surface of sphere of radius r .
 - Since we can choose any shape of the surface enclosing the charge, we choose the simplest possible one! 😊
- The surface is symmetric about the charge.
 - What does this tell us about the field E ?
 - Must have the same magnitude (uniform) at any point on the surface
 - Points radially outward parallel to the surface vector $d\mathbf{A}$.

- The Gaussian integral can be written as

$$\oint \vec{E} \cdot d\vec{A} = \oint E dA = E \oint dA = E(4\pi r^2) = \frac{Q_{encl}}{\epsilon_0} = \frac{Q}{\epsilon_0}$$

Solve
for E

$$E = \frac{Q}{4\pi\epsilon_0 r^2}$$

Electric Field of
Coulomb's Law