

# PHYS 1441 – Section 001

## Lecture #4

*Thursday, June 5, 2014*

*Dr. **Jaehoon** Yu*

- Chapter 2:
  - One Dimensional Motion
    - Acceleration
    - Motion under constant acceleration
    - One dimensional Kinematic Equations
    - How do we solve kinematic problems?
    - Falling motions

Today's homework is homework #2, due 11pm, Tuesday, June 10!!

Thursday, June 5, 2014



PHYS 1441-001, Summer 2014  
Dr. Jaehoon Yu

# Announcements

- Reading Assignment
  - CH2.8
- First term exam
  - In the class coming Monday, June 9
  - Covers CH1.1 through what we learn today (CH2.8?) + Appendix A
  - Bring your calculator but DO NOT input formula into it!
  - You can prepare a one 8.5x11.5 sheet (front and back) of handwritten formulae and values of constants for the exam → no solutions, derivations or definitions!
    - No additional formulae or values of constants will be provided!
- Quiz Results
  - Class average: 50/72
    - Equivalent to 69/100
  - Top score: 72/72



# Displacement, Velocity and Speed

Displacement

$$\Delta x \equiv x_f - x_i$$

Average velocity

$$v_x \equiv \frac{x_f - x_i}{t_f - t_i} = \frac{\Delta x}{\Delta t}$$

Average speed

$$v \equiv \frac{\text{Total Distance Traveled}}{\text{Total Time Spent}}$$

Instantaneous velocity

$$v_x = \lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t} = \frac{dx}{dt}$$

Instantaneous speed

$$|v_x| = \left| \lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t} \right| = \left| \frac{dx}{dt} \right|$$



# Acceleration

Change of velocity in time (what kind of quantity is this?)

• Definition of Average acceleration:

Vector!

$$a_x \equiv \frac{v_{xf} - v_{xi}}{t_f - t_i} = \frac{\Delta v_x}{\Delta t} \quad \text{analogous to} \quad v_x \equiv \frac{x_f - x_i}{t_f - t_i} = \frac{\Delta x}{\Delta t}$$

Dimension?

[LT<sup>-2</sup>]

Unit?


m/s<sup>2</sup>

• Definition of Instantaneous acceleration: Average acceleration over a very short amount of time.

$$a_x \equiv \lim_{\Delta t \rightarrow 0} \frac{\Delta v_x}{\Delta t} \quad \text{analogous to}$$


$$v_x \equiv \lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t}$$

# Acceleration vs Time Plot



# Example 2.4

A car accelerates along a straight road from rest to 75km/h in 5.0s.



What is the magnitude of its average acceleration?

$$v_{xi} = 0 \text{ m/s}$$

$$v_{xf} = \frac{75000 \text{ m}}{3600 \text{ s}} = 21 \text{ m/s}$$

$$a_x = \frac{v_{xf} - v_{xi}}{t_f - t_i} = \frac{\Delta v_x}{\Delta t} = \frac{21 - 0}{5.0} = \frac{21}{5.0} = 4.2 (\text{m/s}^2)$$

$$= \frac{4.2 \times (3600)^2}{1000} = 5.4 \times 10^4 (\text{km/h}^2)$$

# Few Confusing Things on Acceleration

- When an object is moving in a constant velocity ( $v=v_0$ ), there is no acceleration ( $a=0$ )
  - Is there any acceleration when an object is not moving?
- When an object is moving faster as time goes on, ( $v=v(t)$ ), acceleration is positive ( $a>0$ ).
  - Incorrect, since the object might be moving in negative direction initially
- When an object is moving slower as time goes on, ( $v=v(t)$ ), acceleration is negative ( $a<0$ )
  - Incorrect, since the object might be moving in negative direction initially
- In all cases, velocity is positive, unless the direction of the movement changes.
  - Incorrect, since the object might be moving in negative direction initially
- Is there acceleration if an object moves in a constant speed but changes direction?

The answer is YES!!



# The Direction (sign) of the Acceleration

- If the velocity **INCREASES**, the acceleration must be in the **SAME** direction as the velocity!!
  - If the positive velocity increases, what sign is the acceleration?
    - Positive!!
  - If the negative velocity increases, what sign is the acceleration?
    - Negative
- If the velocity **DECREASES**, the acceleration must be in the **OPPOSITE** direction to the velocity!!
  - If the positive velocity decreases, what sign is the acceleration?
    - Negative
  - If the negative velocity decreases, what sign is the acceleration?
    - Positive





# One Dimensional Motion

- Let's focus on the simplest case: acceleration is a constant ( $a=a_0$ )
- Using the definitions of average acceleration and velocity, we can derive equations of motion (description of motion, velocity and position as a function of time)

$$\bar{a}_x = a_x = \frac{v_{xf} - v_{xi}}{t_f - t_i} \quad (\text{If } t_f=t \text{ and } t_i=0) \quad a_x = \frac{v_{xf} - v_{xi}}{t} \quad \Rightarrow \quad v_{xf} = v_{xi} + a_x t$$

For constant acceleration, average velocity is a simple numeric average


$$\bar{v}_x = \frac{v_{xi} + v_{xf}}{2} = \frac{2v_{xi} + a_x t}{2} = v_{xi} + \frac{1}{2} a_x t$$


$$\bar{v}_x = \frac{x_f - x_i}{t_f - t_i} \quad (\text{If } t_f=t \text{ and } t_i=0) \quad \bar{v}_x = \frac{x_f - x_i}{t} \quad \Rightarrow \quad x_f = x_i + \bar{v}_x t$$

Resulting Equation of Motion becomes

$$x_f = x_i + \bar{v}_x t = x_i + v_{xi} t + \frac{1}{2} a_x t^2$$

# One Dimensional Motion cont'd

Average velocity  $\bar{v}_x = \frac{v_{xi} + v_{xf}}{2}$    $x_f = x_i + \bar{v}_x t = x_i + \left( \frac{v_{xi} + v_{xf}}{2} \right) t$

Since  $a_x = \frac{v_{xf} - v_{xi}}{t}$    $t = \frac{v_{xf} - v_{xi}}{a_x}$

Substituting t in the above equation,

$$x_f = x_i + \left( \frac{v_{xf} + v_{xi}}{2} \right) \left( \frac{v_{xf} - v_{xi}}{a_x} \right) = x_i + \frac{v_{xf}^2 - v_{xi}^2}{2a_x}$$

Resulting in

$$v_{xf}^2 = v_{xi}^2 + 2a_x (x_f - x_i)$$

# Kinematic Equations of Motion on a Straight Line Under Constant Acceleration

→  $v_{xf}(t) = v_{xi} + a_x t$  Velocity as a function of time

$$x_f - x_i = \bar{v}_x t = \frac{1}{2} (v_{xf} + v_{xi}) t$$

Displacement as a function of velocities and time

→  $x_f = x_i + v_{xi} t + \frac{1}{2} a_x t^2$  Displacement as a function of time, velocity, and acceleration

→  $v_{xf}^2 = v_{xi}^2 + 2a_x(x_f - x_i)$  Velocity as a function of Displacement and acceleration

You may use different forms of Kinetic equations, depending on the information given to you for specific physical problems!!

# How do we solve a problem using the kinematic formula for constant acceleration?

- Identify what information is given in the problem.
  - Initial and final velocity?
  - Acceleration?
  - Distance?
  - Time?
- Identify what the problem wants you to figure out.
- Identify which kinematic formula is most appropriate and easiest to solve for what the problem wants.
  - Often multiple formulae can give you the answer for the quantity you are looking for. → Do not just use any formula but use the one that makes the problem easiest to solve.
- Solve the equation for the quantity wanted!



# Example

Suppose you want to design an air-bag system that can protect the driver in a head-on collision at a speed 100km/hr (~60miles/hr). Estimate how fast the air-bag must inflate to effectively protect the driver. Assume the car crumples upon impact over a distance of about 1m. How does the use of a seat belt help the driver?

How long does it take for the car to come to a full stop?  As long as it takes for it to crumple.

The initial speed of the car is  $v_{xi} = 100km / h = \frac{100000m}{3600s} = 28m / s$

We also know that  $v_{xf} = 0m / s$  and  $x_f - x_i = 1m$

Using the kinematic formula  $v_{xf}^2 = v_{xi}^2 + 2a_x(x_f - x_i)$

The acceleration is  $a_x = \frac{v_{xf}^2 - v_{xi}^2}{2(x_f - x_i)} = \frac{0 - (28m / s)^2}{2 \times 1m} = -390m / s^2$

Thus the time for air-bag to deploy is  $t = \frac{v_{xf} - v_{xi}}{a} = \frac{0 - 28m / s}{-390m / s^2} = 0.07s$

# Falling Motion

- Falling motion is a motion under the influence of the gravitational pull (gravity) only; Which direction is a freely falling object moving? **Yes, down to the center of the earth!!**
  - A motion under constant acceleration
  - All kinematic formula we learned can be used to solve for falling motions.
- Gravitational acceleration is inversely proportional to the square of the distance between the object and the center of the earth
- The magnitude of the gravitational acceleration is  $g=9.80\text{m/s}^2$  on the surface of the earth.
- The direction of gravitational acceleration is **ALWAYS** toward the center of the earth, which we normally call  $(-y)$ ; where up and down direction are indicated as the variable “y”
- Thus the correct denotation of gravitational acceleration on the surface of the earth is  $g=-9.80\text{m/s}^2$  when  $+y$  points upward
- The difference is that the object initially moving upward will turn around and come down!



# Example for Using 1D Kinematic Equations on a Falling object


A stone was thrown straight upward at  $t=0$  with  $+20.0\text{m/s}$  initial velocity on the roof of a  $50.0\text{m}$  high building,

What is the acceleration in this motion?  $a=g=-9.80\text{m/s}^2$

(a) Find the time the stone reaches at the maximum height.

What happens at the maximum height? The stone stops;  $V=0$

$$v_f = v_{yi} + a_y t = +20.0 - 9.80t = 0.00\text{m/s}$$



$$t = \frac{20.0}{9.80} = 2.04\text{s}$$

(b) Find the maximum height.

$$\begin{aligned} y_f &= y_i + v_{yi}t + \frac{1}{2}a_y t^2 = 50.0 + 20 \times 2.04 + \frac{1}{2} \times (-9.80) \times (2.04)^2 \\ &= 50.0 + 20.4 = 70.4(\text{m}) \end{aligned}$$

# Example of a Falling Object cnt'd

(c) Find the time the stone reaches back to its original height.

$$t = 2.04 \times 2 = 4.08s$$

(d) Find the velocity of the stone when it reaches its original height.

$$v_{yf} = v_{yi} + a_y t = 20.0 + (-9.80) \times 4.08 = -20.0(m / s)$$

(e) Find the velocity and position of the stone at  $t=5.00s$ .

Velocity

$$v_{yf} = v_{yi} + a_y t = 20.0 + (-9.80) \times 5.00 = -29.0(m / s)$$

Position

$$\begin{aligned} y_f &= y_i + v_{yi} t + \frac{1}{2} a_y t^2 \\ &= 50.0 + 20.0 \times 5.00 + \frac{1}{2} \times (-9.80) \times (5.00)^2 = +27.5(m) \end{aligned}$$

