

PHYS 1443 – Section 003

Lecture #11

Monday, Oct. 21, 2002

Dr. Jaehoon Yu

1. Collisions in Two Dimension
2. Center of Mass
 - Definition
 - CM of a Rigid Object
 - Center of Mass and Center of Gravity
3. Motion of a Group of Particles
4. Rocket Propulsion
5. Fundamentals on Rotation

Today's homework is homework #12, due 12:00pm, next Monday!!




Announcements

- 2nd Term exam
 - **Wednesday, Oct. 30**, in the class
 - Covers **chapters 6 – 10**
 - Mixture of Multiple choice and Essay problems
 - Review on Monday, Oct. 28
- Magda Cortez, please come and talk to me after the class



Two dimensional Collisions

In two dimension, one can use components of momentum to apply momentum conservation to solve physical problems.



$$\vec{m}_1 \vec{v}_{1i} + m_2 \vec{v}_{2i} = m_1 \vec{v}_{1f} + m_2 \vec{v}_{2f}$$

x-comp.

$$m_1 v_{1ix} + m_2 v_{2ix} = m_1 v_{1fx} + m_2 v_{2fx}$$

y-comp.

$$m_1 v_{1iy} + m_2 v_{2iy} = m_1 v_{1fy} + m_2 v_{2fy}$$

Consider a system of two particle collisions and scatters in two dimension as shown in the picture. (This is the case at fixed target accelerator experiments.) The momentum conservation tells us:

$$m_1 \vec{v}_{1f} + m_2 \vec{v}_{2f} = m_1 \vec{v}_{1i}$$

$$m_1 v_{1ix} = m_1 v_{1fx} + m_2 v_{2fx} = m_1 v_{1f} \cos q + m_2 v_{2f} \cos f$$

$$m_1 v_{1iy} = 0 = m_1 v_{1fy} + m_2 v_{2fy} = m_1 v_{1f} \sin q - m_2 v_{2f} \sin f$$

And for the elastic conservation, the kinetic energy is conserved:

Monday, Oct. 21, 2002



$$\frac{1}{2} m_1 v_{1i}^2 = \frac{1}{2} m_1 v_{1f}^2 + \frac{1}{2} m_2 v_{2f}^2$$

PHYS 1443-003, Fall 2002
Dr. Jaehoon Yu


What do you think we can learn from these relationships?



Ex09-09.ip

Example 9.9

A 1500kg car traveling east with a speed of 25.0 m/s collides at an intersection with a 2500kg van traveling north at a speed of 20.0 m/s. After the collision the two cars stuck to each other, and the wreckage is moving together. Determine the velocity of the wreckage after the collision, assuming the vehicles underwent a perfectly inelastic collision.



The initial momentum of the two car system before the collision is

$$\begin{aligned}\vec{p}_i &= m_1 v_{1i} \vec{i} + m_2 v_{2i} \vec{j} = 1500 \times 25.0 \vec{i} + 2500 \times 20.0 \vec{j} \\ &= 3.75 \times 10^4 \vec{i} + 5.0 \times 10^4 \vec{j}\end{aligned}$$

The final momentum of the two car system after the perfectly inelastic collision is

$$\vec{p}_f = (m_1 + m_2)(v_{fx} \vec{i} + v_{fy} \vec{j}) = 4.0 \times 10^3 v_{fx} \vec{i} + 4.0 \times 10^3 v_{fy} \vec{j}$$

Using
momentum
conservation

$$\vec{p}_f = \vec{p}_i$$

X-comp.

$$p_{fx} = p_{ix}$$

$$(m_1 + m_2)v_{fx} = m_1 v_{1x} + 0$$

$$v_{fx} = \frac{(m_1 v_{1x} + 0)}{m_1 + m_2} = \frac{3.75 \times 10^4}{1500 + 2500} = 9.38 \text{ m/s}$$

Y-comp.

$$p_{fy} = p_{iy}$$

$$(m_1 + m_2)v_{fy} = 0 + m_2 v_{2y}$$

$$v_{fy} = \frac{(0 + m_2 v_{2y})}{m_1 + m_2} = \frac{5.0 \times 10^4}{1500 + 2500} = 12.5 \text{ m/s}$$

$$\vec{v}_f = v_{fx} \vec{i} + v_{fy} \vec{j} = (9.38 \vec{i} + 12.5 \vec{j}) \text{ m/s}$$

Monday, Oct. 21, 2002




PHYS 1443-003, Fall 2002

Dr. Jaehoon Yu

Example 9.10

Proton #1 with a speed 3.50×10^5 m/s collides elastically with proton #2 initially at rest. After the collision, proton #1 moves at an angle of 37° to the horizontal axis and proton #2 deflects at an angle ϕ to the same axis. Find the final speeds of the two protons and the scattering angle of proton #2, ϕ .



From kinetic energy conservation:

$$(3.50 \times 10^5)^2 = v_{1f}^2 + v_{2f}^2 \quad (3)$$

Since both the particles are protons $m_1 = m_2 = m_p$.

Using momentum conservation, one obtains

x-comp. $m_p v_{1i} = m_p v_{1f} \cos q + m_p v_{2f} \cos f$

y-comp. $m_p v_{1f} \sin q - m_p v_{2f} \sin f = 0$

Canceling m_p and put in all known quantities, one obtains

$$v_{1f} \cos 37^\circ + v_{2f} \cos f = 3.50 \times 10^5 \quad (1)$$

$$v_{1f} \sin 37^\circ = v_{2f} \sin f \quad (2)$$

Solving Eqs. 1-3 equations, one gets

$$v_{1f} = 2.80 \times 10^5 \text{ m / s}$$

$$v_{2f} = 2.11 \times 10^5 \text{ m / s}$$

$$f = 53.0^\circ$$


Center of Mass

We've been solving physical problems treating objects as sizeless points with masses, but in realistic situation objects have shapes with masses distributed throughout the body.

Center of mass of a system is the average position of the system's mass and represents the motion of the system as if all the mass is on the point.

What does above statement tell you concerning forces being exerted on the system?

The total external force exerted on the system of total mass M causes the center of mass to move at an acceleration given by $\vec{a} = \sum \vec{F} / M$ as if all the mass of the system is concentrated on the center of mass.



Consider a massless rod with two balls attached at either end.

The position of the center of mass of this system is the mass averaged position of the system

$$x_{CM} \equiv \frac{m_1 x_1 + m_2 x_2}{m_1 + m_2}$$

CM is closer to the heavier object



Center of Mass of a Rigid Object

The formula for CM can be expanded to Rigid Object or a system of many particles

$$x_{CM} = \frac{m_1 x_1 + m_2 x_2 + \cdots + m_n x_n}{m_1 + m_2 + \cdots + m_n} = \frac{\sum_i m_i x_i}{\sum_i m_i}$$


$$y_{CM} = \frac{\sum_i m_i y_i}{\sum_i m_i}$$

$$z_{CM} = \frac{\sum_i m_i z_i}{\sum_i m_i}$$

The position vector of the center of mass of a many particle system is

$$\vec{r}_{CM} = x_{CM} \vec{i} + y_{CM} \vec{j} + z_{CM} \vec{k} = \frac{\sum_i m_i x_i \vec{i} + \sum_i m_i y_i \vec{j} + \sum_i m_i z_i \vec{k}}{\sum_i m_i}$$

$$\vec{r}_{CM} = \frac{\sum_i m_i \vec{r}_i}{M}$$



A rigid body – an object with shape and size with mass spread throughout the body, ordinary objects – can be considered as a group of particles with mass m_i densely spread throughout the given shape of the object

$$x_{CM} \approx \frac{\sum_i \Delta m_i x_i}{M}$$


$$x_{CM} = \lim_{\Delta m_i \rightarrow 0} \frac{\sum_i \Delta m_i x_i}{M} = \frac{1}{M} \int x dm$$

$$\vec{r}_{CM} = \frac{1}{M} \int \vec{r} dm$$



Center of Mass and Center of Gravity

The center of mass of any symmetric object lies on an axis of symmetry and on any plane of symmetry, if object's mass is evenly distributed throughout the body.




How do you think you can determine the CM of objects that are not symmetric?

One can use gravity to locate CM.

1. Hang the object by one point and draw a vertical line following a plum-bob.
2. Hang the object by another point and do the same.
3. The point where the two lines meet is the CM.

Center of Gravity



Since a rigid object can be considered as collection of small masses, one can see the total gravitational force exerted on the object as


$$\vec{F}_g = \sum_i \vec{F}_i = \sum_i \Delta m_i \vec{g} = M \vec{g}$$

What does this equation tell you?

The net effect of these small gravitational forces is equivalent to a single force acting on a point (**Center of Gravity**) with mass M.

Example 9.12

A system consists of three particles as shown in the figure. Find the position of the center of mass of this system.



Using the formula for CM for each position vector component

$$x_{CM} = \frac{\sum m_i x_i}{\sum m_i}$$

$$y_{CM} = \frac{\sum m_i y_i}{\sum m_i}$$

One obtains $\vec{r}_{CM} = x_{CM} \vec{i} + y_{CM} \vec{j} = \frac{(m_2 + 2m_3) \vec{i} + 2m_1 \vec{j}}{m_1 + m_2 + m_3}$

$$x_{CM} = \frac{\sum m_i x_i}{\sum m_i} = \frac{m_1 x_1 + m_2 x_2 + m_3 x_3}{m_1 + m_2 + m_3} = \frac{m_2 + 2m_3}{m_1 + m_2 + m_3}$$

$$y_{CM} = \frac{\sum m_i y_i}{\sum m_i} = \frac{m_1 y_1 + m_2 y_2 + m_3 y_3}{m_1 + m_2 + m_3} = \frac{2m_1}{m_1 + m_2 + m_3}$$


If $m_1 = 2\text{kg}; m_2 = m_3 = 1\text{kg}$

$$\vec{r}_{CM} = \frac{3\vec{i} + 4\vec{j}}{4} = 0.75\vec{i} + \vec{j}$$



Example 9.13

Show that the center of mass of a rod of mass M and length L lies in midway between its ends, assuming the rod has a uniform mass per unit length.



The formula for CM of a continuous object is

$$x_{CM} = \frac{1}{M} \int_{x=0}^{x=L} x dm$$

Since the density of the rod (λ) is constant; $\lambda = M/L$

The mass of a small segment $dm = \lambda dx$

Therefore

$$x_{CM} = \frac{1}{M} \int_{x=0}^{x=L} \lambda x dx = \frac{1}{M} \left[\frac{1}{2} \lambda x^2 \right]_{x=0}^{x=L} = \frac{1}{M} \left(\frac{1}{2} \lambda L^2 \right) = \frac{1}{M} \left(\frac{1}{2} ML \right) = \frac{L}{2}$$

Find the CM when the density of the rod non-uniform but varies linearly as a function of x , $\lambda = \alpha x$

$$M = \int_{x=0}^{x=L} \lambda dx = \int_{x=0}^{x=L} \alpha x dx$$

$$= \left[\frac{1}{2} \alpha x^2 \right]_{x=0}^{x=L} = \frac{1}{2} \alpha L^2$$

Monday, Oct. 21, 2002



PHYS 144C
Dr. Jaehoon Yu

$$x_{CM} = \frac{1}{M} \int_{x=0}^{x=L} \lambda x dx = \frac{1}{M} \int_{x=0}^{x=L} \alpha x^2 dx = \frac{1}{M} \left[\frac{1}{3} \alpha x^3 \right]_{x=0}^{x=L}$$

$$x_{CM} = \frac{1}{M} \left(\frac{1}{3} \alpha L^3 \right) = \frac{1}{M} \left(\frac{2}{3} ML \right) = \frac{2L}{3}$$

Motion of a Group of Particles

We've learned that the CM of a system can represent the motion of a system. Therefore, for an isolated system of many particles in which the total mass M is preserved, the velocity, total momentum, acceleration of the system are

Velocity of the system

$$\vec{v}_{CM} = \frac{d\vec{r}_{CM}}{dt} = \frac{d}{dt} \left(\frac{1}{M} \sum m_i \vec{r}_i \right) = \frac{1}{M} \sum m_i \frac{d\vec{r}_i}{dt} = \frac{\sum m_i \vec{v}_i}{M}$$

Total Momentum of the system

$$\vec{p}_{CM} = M \vec{v}_{CM} = M \frac{\sum m_i \vec{v}_i}{M} = \sum m_i \vec{v}_i = \sum \vec{p}_i = \vec{p}_{tot}$$

Acceleration of the system

$$\vec{a}_{CM} = \frac{d\vec{v}_{CM}}{dt} = \frac{d}{dt} \left(\frac{1}{M} \sum m_i \vec{v}_i \right) = \frac{1}{M} \sum m_i \frac{d\vec{v}_i}{dt} = \frac{\sum m_i \vec{a}_i}{M}$$

External force exerting on the system

$$\sum \vec{F}_{ext} = M \vec{a}_{CM} = \sum m_i \vec{a}_i = \frac{d\vec{p}_{tot}}{dt}$$

What about the internal forces?

If net external force is 0

$$\sum \vec{F}_{ext} = 0 = \frac{d\vec{p}_{tot}}{dt}$$

$$\vec{p}_{tot} = \text{const}$$

System's momentum is conserved.

Monday, Oct. 21, 2002



PHYS 1443-003, Fall 2002
Dr. Jaehoon Yu

11