PHYS 1443 – Section 003 Lecture #17

Wednesday, Oct. 27, 2004 Dr. **Jae**hoon Yu

- 1. Fundamentals on Rotational Motion
- 2. Rotational Kinematics
- 3. Relationship between angular and linear quantities
- 4. Rolling Motion of a Rigid Body
- 5. Torque

2nd Term Exam Monday, Nov. 1!! Covers CH 6 – 10.5!!

No homework today!!



Fundamentals on Rotation

Linear motions can be described as the motion of the center of mass with all the mass of the object concentrated on it.

Is this still true for rotational motions?



No, because different parts of the object have different linear velocities and accelerations.

Consider a motion of a rigid body – an object that does not change its shape – rotating about the axis protruding out of the slide.

The arc length, or sergita, is $l = R \theta$

Therefore the angle, θ , is $\theta = \frac{l}{R}$. And the unit of the angle is in radian. It is dimensionless!!

One radian is the angle swept by an arc length equal to the radius of the arc.

Since the circumference of a circle is $2\pi r$, $360^{\circ} = 2\pi r / r = 2\pi$

The relationship between radian and degrees is $1 \operatorname{rad} = 360^{\circ} / 2\pi = 180^{\circ} / \pi$

Wednesday, Oct. 27, 2004



PHYS 1443-003, Fall 2004 Dr. Jaehoon Yu $\approx 180^{\circ}/3.14 \approx 57.3^{\circ}$ ²

Example 10 – 1

A particular bird's eyes can barely distinguish objects that subtend an angle no smaller than about $3x10^{-4}$ rad. (a) How many degrees is this? (b) How small an object can the bird just distinguish when flying at a height of 100m?



(a) One radian is $360^{\circ}/2\pi$. Thus (b) $3 \times 10^{-4} rad = (3 \times 10^{-4} rad) \times$ $(360^{\circ}/2\pi \, rad) = 0.017^{\circ}$ (b) Since $I=r\theta$ and for small angle arc length is approximately the same as the chord length. $l = r\theta =$ Chord $100m \times 3 \times 10^{-4} rad =$ $3 \times 10^{-2} m = 3 cm$ Arc length 3

Rotational Kinematics

The first type of motion we have learned in linear kinematics was under a constant acceleration. We will learn about the rotational motion under constant angular acceleration, because these are the simplest motions in both cases.

Just like the case in linear motion, one can obtain

Angular Speed under constant angular acceleration:

Angular displacement under constant angular acceleration:

One can also obtain

$$\omega_{f} = \omega_{i} + \alpha t$$

$$\theta_{f} = \theta_{i} + \omega_{i}t + \frac{1}{2}\alpha t^{2}$$

$$\omega_{f}^{2} = \omega_{i}^{2} + 2\alpha \left(\theta_{f} - \theta_{i}\right)$$



Angular Displacement, Velocity, and Acceleration

Using what we have learned in the previous slide, how would you define the angular displacement?

- How about the average angular speed? Unit? rad/s
- And the instantaneous angular speed? Unit? rad/s

By the same token, the average angular acceleration Unit? rad/s² And the instantaneous angular acceleration? Unit? rad/s²

$$\overline{\omega} = \frac{\theta_f - \theta_i}{t_f - t_i} = \frac{\Delta \theta}{\Delta t}$$

$$\omega \equiv \lim_{\Delta t \to 0} \frac{\Delta \theta}{\Delta t} = \frac{d\theta}{dt}$$

$$\overline{\alpha} \equiv \frac{\omega_f - \omega_i}{t_f - t_i} = \frac{\Delta \omega}{\Delta t}$$

$$\alpha \equiv \lim_{\Delta t \to 0} \frac{\Delta \omega}{\Delta t} = \frac{d\omega}{dt}$$

 $\Delta \theta = \theta_f - \theta_i$

When rotating about a fixed axis, every particle on a rigid object rotates through the same angle and has the same angular speed and angular acceleration.



Example for Rotational Kinematics

A wheel rotates with a constant angular acceleration of 3.50 rad/s². If the angular speed of the wheel is 2.00 rad/s at $t_i=0$, a) through what angle does the wheel rotate in 2.00s?

Using the angular displacement formula in the previous slide, one gets

$$\begin{aligned} \theta_f - \theta_i &= \omega t + \frac{1}{2} \alpha t^2 \\ = 2.00 \times 2.00 + \frac{1}{2} 3.50 \times (2.00)^2 &= 11.0 \, rad \\ &= \frac{11.0}{2\pi} \, rev. = 1.75 \, rev. \end{aligned}$$



Example for Rotational Kinematics cnt'd

What is the angular speed at t=2.00s?

Using the angular speed and acceleration relationship

$$\omega_f = \omega_i + \alpha t = 2.00 + 3.50 \times 2.00 = 9.00 rad/s$$

Find the angle through which the wheel rotates between t=2.00 s and t=3.00 s.

Using the angular kinematic formula $\theta_f - \theta_i = \omega t + \frac{1}{2} \alpha t^2$ At t=2.00s $\theta_{t=2} = 2.00 \times 2.00 + \frac{1}{2} 3.50 \times 2.00 = 11.0 rad$ At t=3.00s $\theta_{t=3} = 2.00 \times 3.00 + \frac{1}{2} 3.50 \times (3.00)^2 = 21.8 rad$ Angular $\Delta \theta = \theta_3 - \theta_2 = 10.8 rad = \frac{10.8}{2\pi} rev. = 1.72 rev.$ We driesday, oct. 27, 2004 Θ PHYS 1443-003, Fall 2004 T



What does this relationship tell you about the tangential speed of the points in the object and their angular speed?:

Wednesday, Oct. 27, 2004



PHYS 1 The farther away the particle is from the center of Di rotation, the higher the tangential speed.

angular speed, its tangential speed differs

Although every particle in the object has the same

proportional to its distance from the axis of rotation.

8

Is the lion faster than the horse?

A rotating carousel has one child sitting on a horse near the outer edge and another child on a lion halfway out from the center. (a) Which child has the greater linear speed? (b) Which child has the greater angular speed?



(a) Linear speed is the distance traveled divided by the time interval. So the child sitting at the outer edge travels more distance within the given time than the child sitting closer to the center. Thus, the horse is faster than the lion.

(b) Angular speed is the angle traveled divided by the time interval. The angle both the children travel in the given time interval is the same. Thus, both the horse and the lion have the same angular speed.





What does this relationship tell you?

How about the acceleration?

How many different linear accelerations do you see in a circular motion and what are they? Two

Tangential, $a_{t'}$ and the radial acceleration, a_r

Since the tangential speed v is $v = r\omega$

The magnitude of tangential acceleration a_t is

$$a_t = \frac{dv}{dt} = \frac{d}{dt}(r\omega) = r\frac{d\omega}{dt} = r\alpha$$

Although every particle in the object has the same angular acceleration, its tangential acceleration differs proportional to its distance from the axis of rotation.

The radial or centripetal acceleration a_r is

$$a_r = \frac{v^2}{r} = \frac{(r\omega)^2}{r} = r\omega^2$$

What doesThe father away the particle is from the rotation axis, the more radialthis tell you?acceleration it receives. In other words, it receives more centripetal force.

Total linear acceleration is

$$a = \sqrt{a_t^2 + a_r^2} = \sqrt{(r\alpha)^2 + (r\omega^2)^2} = r\sqrt{\alpha^2 + \omega^4}$$

Wednesday, Oct. 27, 2004



PHYS 1443-003, Fall 2004 Dr. Jaehoon Yu

Example

(a) What is the linear speed of a child seated 1.2m from the center of a steadily rotating merry-go-around that makes one complete revolution in 4.0s? (b) What is her total linear acceleration?

First, figure out what the angular speed of the merry-go-around is.

$$\varpi = \frac{1rev}{4.0s} = \frac{2\pi rad}{4.0s} = 1.6rad / s$$

Using the formula for linear speed

$$v = r\omega = 1.2m \times 1.6rad / s = 1.9m / s$$

Since the angular speed is constant, there is no angular acceleration.

Tangential acceleration is

Radial acceleration is

Thus the total acceleration is

$$a_{t} = r\alpha = 1.2m \times 0 rad / s^{2} = 0m / s^{2}$$

$$a_{r} = r\varpi^{2} = 1.2m \times (1.6rad / s)^{2} = 3.1m / s^{2}$$

$$a_{r} = \sqrt{a_{t}^{2} + a_{r}^{2}} = \sqrt{0 + (3.1)^{2}} = 3.1m / s^{2}$$



Example for Rotational Motion

Audio information on compact discs are transmitted digitally through the readout system consisting of laser and lenses. The digital information on the disc are stored by the pits and flat areas on the track. Since the speed of readout system is constant, it reads out the same number of pits and flats in the same time interval. In other words, the linear speed is the same no matter which track is played. a) Assuming the linear speed is 1.3 m/s, find the angular speed of the disc in revolutions per minute when the inner most (r=23mm) and outer most tracks (r=58mm) are read.

Using the relationship
between angular and
tangential speed
$$v = r\omega$$

b) The maximum playing time of a standard music
CD is 74 minutes and 33 seconds. How many
revolutions does the disk make during that time?
c) What is the total length of the track past through the readout mechanism?
c) What is the angular acceleration of the CD over
the 4473s time interval, assuming constant α ?
 $\alpha = \frac{(\omega_f - \omega_i)}{\Delta t} = \frac{(22.4 - 56.5)rad/s}{4473s}$
Wednesday, Oct. 27, 200
 $\alpha = \frac{(\omega_f - \omega_i)}{Dr. Jaehoon Yu}$
 $\alpha = 7.6 \times 10^{-3} rad/s^2$ 12

Rolling Motion of a Rigid Body

What is a rolling motion?

A more generalized case of a motion where the rotational axis moves together with the object

To simplify the discussion, let's make a few assumptions

A rotational motion about the moving axis

- 1. Limit our discussion on very symmetric objects, such as cylinders, spheres, etc
- 2. The object rolls on a flat surface

Let's consider a cylinder rolling without slipping on a flat surface

R S The to

Under what condition does this "Pure Rolling" happen?

The total linear distance the CM of the cylinder moved is

$$s = R\theta$$

Thus the linear speed of the CM is

$$v_{CM} = \frac{ds}{dt} = R\frac{d\theta}{dt} = R\omega$$

Wednesday, Oct. 27, 2004



PHYS 1443-003, Fall 2004 Condition for "Pure Rolling" 13 Dr. Jaehoon Yu

More Rolling Motion of a Rigid Body

The magnitude of the linear acceleration of the CM is

$$a_{CM} = \frac{dv_{CM}}{dt} = R\frac{d\omega}{dt} = R\alpha$$

Why??



As we learned in the rotational motion, all points in a rigid body moves at the same angular speed but at a different linear speed. CM is moving at the same speed at all times.

At any given time, the point that comes to P has 0 linear speed while the point at P' has twice the speed of CM

A rolling motion can be interpreted as the sum of Translation and Rotation



Torque

Torque is the tendency of a force to rotate an object about an axis. Torque, τ , is a vector quantity.



Consider an object pivoting about the point P by the force \mathcal{F} being exerted at a distance r. The line that extends out of the tail of the force vector is called the line of action.

The perpendicular distance from the pivoting point P to the line of action is called Moment arm.

Magnitude of torque is defined as the product of the force exerted on the object to rotate it and the moment arm.

When there are more than one force being exerted on certain points of the object, one can sum up the torque generated by each force vectorially. The convention for sign of the torque is positive if rotation is in counter-clockwise and negative if clockwise.

Wednesday, Oct. 27, 2004



PHYS 1443-003, Fall 2004 Dr. Jaehoon Yu

$$\tau \equiv rF\sin\phi = Fd$$

$$\sum \tau = \tau_1 + \tau_2$$

$$=F_1d_1-F_2d_2$$

Example for Torque

A one piece cylinder is shaped as in the figure with core section protruding from the larger drum. The cylinder is free to rotate around the central axis shown in the picture. A rope wrapped around the drum whose radius is \mathcal{R}_1 exerts force \mathcal{F}_1 to the right on the cylinder, and another force exerts \mathcal{F}_2 on the core whose radius is \mathcal{R}_2 downward on the cylinder. A) What is the net torque acting on the cylinder about the rotation axis?



The torque due to
$$\mathbf{F}_1$$
 $\tau_1 = -R_1 F_1$ and due to \mathbf{F}_2 $\tau_2 = R_2 F_2$

So the total torque acting on the system by the forces is

$$\sum \tau = \tau_1 + \tau_2 = -R_1 F_1 + R_2 F_2$$

Suppose $F_1=5.0$ N, $R_1=1.0$ m, $F_2=15.0$ N, and $R_2=0.50$ m. What is the net torque about the rotation axis and which way does the cylinder rotate from the rest?

Using the above result

$$\sum \tau = -R_1 F_1 + R_2 F_2$$

= -5.0×1.0+15.0×0.50=2.5N•m The cylinder rotates in Counter-clockwise.

